

Sagittal balance of the spine - lumbar lordosis or lumbosacral lordosis?

Kai Song¹✉, Pengfei Chi¹, Qiang Yang², Cao Yang³, Bing Wang⁴, Fangcai Li⁵, Zezhang Zhu⁶, Weishi Li⁷, Jianguo Zhang⁸, Zheng Wang^{1*}✉

1. Department of Orthopaedics, Chinese People's Liberation Army General Hospital, Beijing, PR China.
2. Department of Spine Surgery, Tianjin Hospital, Tianjin, PR China
3. Department of Orthopaedics, Union Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, Hubei, PR China
4. Department of Spine Surgery, Second Xiangya Hospital of Central South University, Changsha, Hunan, PR China
5. Department of Orthopedics Surgery, Second Affiliated Hospital, School of Medicine, Zhejiang University, Hangzhou, Zhejiang, PR China
6. Department of Spine Surgery, The Affiliated Drum Tower Hospital of Nanjing University Medical School, Nanjing, Jiangsu, PR China
7. Department of Orthopaedics, Peking University Third Hospital, Beijing, PR China
8. Department of Orthopedics, Peking Union Medical College Hospital, Beijing, PR China
9. Shenzhen Institute of Advanced Technology, Chinese Academy of Sciences, Shenzhen, PR China

Kai Song ✉: songkai200018@163.com

Wang Zheng* ✉: wangzheng301@163.com

Abstract

Objective: To investigate sagittal spinal alignment from the perspective of the overall curvature of the "S" curve of the human spine, and explore the roles of pelvic incidence (PI) and maximal thoracolumbar vertebral tilt (TLmax) in the classification of the sagittal spinal alignment.

Methods: The tilt of the sacral, lumbar, and thoracic vertebrae (from Co1, S5, S4... to C7) were measured. The minimal sacral vertebral tilt (Smin), maximal thoracolumbar vertebral tilt (TLmax), and minimal thoracic vertebral tilt (Tmin) were recorded. The concept of lumbosacral lordosis (LSL) was introduced, and the Ferguson method was utilized to measure sagittal spinal parameters both in anatomical segmentation (Ferguson L1-S2, Ferguson T1-T12) and functional segmentation (Ferguson LSLmax, Ferguson TKmax). The subjects were grouped based on pelvic incidence (PI) and TLmax separately, and the mean and standard deviation of each parameter were calculated. Chi-square tests were conducted for statistical analysis.

Results: 1. Based on PI grouping: PI for all subjects was $45.4 \pm 9.5^\circ$ (21.7-86.4°). Group A consisted of 117 subjects with a mean PI of $34.7 \pm 4.4^\circ$, Group B had 158 subjects with a mean PI of $45.2 \pm 2.9^\circ$, and Group C included 113 subjects with a mean PI of $56.7 \pm 5.8^\circ$. No statistically significant differences were found in tilt of S2, L1, T1, TLmax, Tmin, and Ferguson L1-S2, Ferguson T1-T12, and Ferguson TKmax among Groups A, B, and C. 2. Based on TLmax grouping: TLmax for all subjects was $110.5 \pm 5.5^\circ$ (94.4-132.0°). Group A had 91 subjects with a mean TLmax of $104.0 \pm 2.3^\circ$, Group B comprised 216 subjects with a mean TLmax of $110.2 \pm 2.1^\circ$, and Group C included 81 subjects with a mean TLmax of $118.6 \pm 3.8^\circ$. Significant statistical differences were observed in tilt of S2, L1, T1, Smin, TLmax, Tmin, and Ferguson L1-S2, Ferguson T1-T12, Ferguson LSLmax, and Ferguson TKmax among Groups A, B, and C.

Conclusion: There were no differences in the magnitude of LSL and TK among subjects with different PI, indicating that PI does not affect the overall curvature of the "S" curve in the sagittal spinal alignment. In contrast, TLmax effectively distinguishes the

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

Keywords: Sagittal spinal alignment; Sagittal spinal curvature; Lumbosacral lordosis; Pelvic incidence; Maximal thoracolumbar vertebral tilt.

Introduction

The recognition of sagittal spinal alignments is of significant importance for the diagnosis and treatment of spinal disorders [1, 2]. The "S" curve formed by the lumbar lordosis and thoracic kyphosis has played a crucial role in the evolutionary process of humans transitioning from quadrupedal to bipedal upright locomotion [3-6]. Typically, scholars define the lumbar lordosis (LL) as the Cobb angle from L1 to S1 and the thoracic kyphosis (TK) as the Cobb angle from T1 to T12, to anatomically segment and analyze the "S" curve in radiographic images [7].

In recent years, there has been a preference among scholars to use functional segmentation based on the curvature of the "S" curve, rather than rigidly using the upper endplate of L1 as the boundary between the two bends. This approach is similar to the concept of the end vertebrae commonly used in scoliosis, where the inflection point's position is not determined by the anatomical features of the vertebrae but by the characteristics of the spinal sequence [8-11]. Consequently, the cephalic end of LL may be located either proximally or distally on L1. In contrast, the caudal starting point of the lordosis is consistently defined by the upper endplate of S1.

Contrary to the previous study that considered the sacrum as a whole kyphotic segment [12-14], Song et al. confirmed that, in terms of spinal sequence, the sacrum is not entirely kyphotic but is divided into a lordotic segment (S1-S2) and a kyphotic segment (S2-S5, Co1). S1 and S2 actually continue the LL, and S2 should be considered the caudal vertebra of the lordotic segment in the sagittal spinal alignment [15]. If this is the case, the traditional definition of the "S" curve is incomplete, as it lacks the sacral lordotic segment at its distal end. In the context of studying spinal sequences, the definition of lumbosacral lordosis (LSL) seems more reasonable than lumbar lordosis alone.

Based on this background, the present study aims to measure and analyze the spinal curvature of the "S" curve in the general population from both anatomical segmentation and functional segmentation, using vertebral tilt and the Ferguson method. The study also considers the maximum thoracolumbar tilt (TLmax) as a reference for classification and typing of the sagittal spinal curvature, comparing it with previous classification types based on pelvic incidence (PI) [9-11, 16-19]. This comparison aims to provide a deeper exploration of the characteristics of the human sagittal spinal alignments and offer references for the diagnosis and treatment of spinal disorders.

Methods

The study included 388 healthy adults aged 18-35 years old and collected their full-length lateral spine X-rays in a natural standing position. Among them, 336 cases were from Chinese PLA General Hospital, where they had been suspected of having scoliosis by other hospitals, but subsequent outpatient imaging examinations confirmed no abnormalities; 52 cases were from collaborating hospitals. Ethics committee of Chinese PLA General Hospital gave ethical approval (S2023-446-02) for this work.

1. Measurements: The tilt (right angle to the horizontal line) of the anterior, middle, and posterior edges of the vertebrae from Co, S5, S4... to C7 were measured. The minimal sacral vertebral tilt (Smin), the maximal thoracolumbar vertebral tilt (TLmax), and the minimal thoracic vertebral tilt (Tmin) were recorded. The lumbosacral lordosis angle (Ferguson L1-S2), thoracic kyphosis angle (Ferguson T1-T12), maximal lumbosacral lordosis angle (Ferguson LSLmax, calculated as Ferguson Smin-TLmax), and maximal thoracic kyphosis angle (Ferguson TKmax, calculated as Ferguson TLmax-Tmin) were calculated.

2. Conventional measurements: Pelvic incidence (PI), sacral slope (SS), pelvic tilt (PT), lumbar lordosis angle (LL, Cobb L1-S1), thoracic kyphosis angle (TK, Cobb T1-T12), T9 spino-pelvic inclination (T9-SPi), T1 spino-pelvic inclination (T1-SPi), T1 pelvic angle (TPA), spine-sacral angle (SSPA), and sagittal-vertical-axis angle (SVA-A) (the angle of the perpendicular line from the posterior edge of the S1 upper endplate to the center point of the C7 vertebra, with the right angle considered positive, thus converting the SVA distance value into an angle value to eliminate height differences) were measured.

3. Subject grouping: Based on pelvic incidence (PI): Group A $<40^\circ$; Group B $40^\circ \leq PI < 50^\circ$; Group C $\geq 50^\circ$. Based on the maximal thoracolumbar vertebral tilt (TLmax) (middle): Group A $<106.5^\circ$; Group B $106.5^\circ \leq TLmax < 114.5^\circ$; Group C $\geq 114.5^\circ$. The mean, standard deviation, and extreme values of each parameter were calculated, and chi-square tests were performed to compare and analyze the groups within the two classification types.

Results

The measured and statistical parameters are presented in **Tables 1, 2, 3, and 4**.

Based on PI grouping: The overall mean PI for all subjects was $45.4 \pm 9.5^\circ$ (range 21.7 - 86.4°). Group A consisted of 117 subjects with a mean PI of $34.7 \pm 4.4^\circ$ (range 21.7 - 40.0°); Group B had 158 subjects with a mean PI of $45.2 \pm 2.9^\circ$ (range 40.0 - 50.0°); Group C included 113 subjects with a mean PI of $56.7 \pm 5.8^\circ$ (range 50.2 - 86.4°). There were no significant statistical differences in the tilt of S2, L1, T8, T6, T5, T4, T3, T2, T1, C7, TLmax, Tmin among Groups A, B, and C; no significant statistical differences were found for Ferguson L1-S2, Ferguson T1-T12, FergusonTKmax; no significant statistical differences for Cobb T1-T12, SVA-A; however, other parameters showed statistical differences. (**Tables 1, 2**)

Based on TLmax grouping: The overall mean TLmax for all subjects was $110.5 \pm 5.5^\circ$ (range 94.4 - 132.0°). Group A had 91 subjects with a mean TLmax of $104.0 \pm 2.3^\circ$ (range 94.4 - 106.5°); Group B comprised 216 subjects with a mean TLmax of $110.2 \pm 2.1^\circ$ (range 106.6 - 114.5°); Group C included 81 subjects with a mean TLmax of $118.6 \pm 3.8^\circ$ (range 114.7 - 132.0°). There were no significant statistical differences in the tilt of Co1, S5, S4, S3, C7 among Groups A, B, and C, nor in PI, SS; a tendency for statistical differences was observed for S2, T8; other parameters showed statistical differences. (**Tables 3, 4**)

Discussion

The evolution of humans from quadrupedal to bipedal locomotion, with the emergence of the "S" curve of the spine, which provided the necessary support for long-term upright activities with lower energy expenditure, facilitating their survival, reproduction, and development[3-6]. The emergence of the anterior convexity of the "S" curve is particularly important. It is commonly believed that the lordotic sequence is composed of the lumbar vertebrae and intervertebral discs, hence, most scholars report the angle between the upper endplates of L1 and S1 as the measurement index for this curve (lumbar lordosis, LL) [7]. However, from the perspective of spinal sequence, even in the normal population, the lordotic region does not strictly adhere to anatomical features, which has led to much confusion and debate in the assessment and application of the sagittal spinal alignment. Therefore, many scholars have redefined the "S" curve based on functional characteristics, no longer rigidly using the upper endplate of L1 as the boundary between the two bends, but instead, defining the boundary based on the inflection point of the spinal curvature. The definition of the inflection point is similar to the concept of the end vertebra commonly used in scoliosis, and its position does not depend on anatomical features. Consequently, the cephalic end of the lumbar lordosis may be located either proximally or distally on L1. Unlike the uncertainty of the cephalic end, the upper endplate of S1 is consistently used as the starting point of the lordosis, which may be related to the unique anatomical features of the sacrum.

However, contrary to previous studies that considered the sacrum as a whole kyphotic segment and S2 as the apex of kyphosis [12-14], Song et al.'s radiographic study demonstrated that, in terms of spinal sequence, the sacrum is not entirely kyphotic but is divided into a lordotic segment (S1-S2) and a kyphotic segment (S2-S5, Co1), with S1 and S2 effectively continuing the LL, and S2 should be considered the caudal end vertebrae of the lordosis [15]. Song et al. argued that, anatomically, S1 and S2 are within the tension band of the lumbar lordosis, as the erector spinae muscles terminate at the lamina and spinous process of S1 and S2, while the sacrotuberous ligament, sacrospinous ligament, and anococcygeal ligament insert at S3, S4, S5, and Co1, and their reverse tension determines the kyphotic characteristics of this region. Moreover, S2 is the stress core and rotational center of the sacroiliac joint, which is more consistent with the mechanical characteristics of an end vertebra rather than an apex vertebra [15]. Therefore, based on the radiographic study results and logical deductions from anatomy and biomechanics, S2 should be considered the caudal vertebra of the lordosis in the sagittal spinal alignment. The previous definition of the "S" curve's lordotic range was incomplete, as it lacked a part of the sacral lordosis at its distal end. Thus, the definition of lumbosacral lordosis (LSL) seems more reasonable than that of lumbar lordosis (LL). The main reason for this omission is likely due to the limitations of the Cobb method in measuring the sacral region. For the lumbar spine, the Cobb method is relatively applicable due to the distinct boundaries provided by the larger intervertebral discs. However, for the sacral region, which undergoes fusion during development and significant morphological changes, making the Cobb method less applicable. This has led to the incompleteness of the radiographic evaluation for lordosis. To further explore the sagittal spinal alignment in humans, this study introduces the measurement using vertebral tilt and the Ferguson method, with a focus on studying LSL rather than LL.

The study compares two classifications of the sagittal spinal alignment in a healthy population: 1) classification based on pelvic incidence (PI); 2) classification based on the maximal thoracolumbar vertebral tilt(TLmax).

It is made available under a [CC-BY-NC 4.0 International license](https://creativecommons.org/licenses/by-nc/4.0/).

Table 1 Classification based on PI - vertebral tilt and Ferguson angle of sagittal spinal alignment

		Total (n=388)			Classification of PI						F	P
					Group A(n=117)		Group B(n=158)		Group C(n=113)			
anterior edge of the vertebral body	PI	45.4 ± 9.5	(21.7 - 86.4)	34.7 ± 4.4	(21.7 - 40.0)	45.2 ± 2.9	(40.0 - 50.0)	56.7 ± 5.8	(50.2 - 86.4)	724.785	0.000	
	Co1	82.0 ± 13.5	(43.2 - 117.7)	75.8 ± 12.2	(43.2 - 106.8)	82.2 ± 12.5	(44.0 - 117.7)	88.1 ± 13.4	(56.3 - 116.6)	26.723	0.000	
	S5	80.4 ± 12.0	(40.7 - 117.7)	72.8 ± 11.8	(40.7 - 99.9)	81.3 ± 10.0	(50.2 - 105.7)	87.2 ± 10.4	(57.0 - 117.7)	52.530	0.000	
	S4	61.6 ± 12.9	(25.7 - 92.9)	53.3 ± 11.0	(30.7 - 84.3)	61.5 ± 11.4	(25.7 - 84.6)	70.3 ± 11.2	(45.0 - 92.9)	65.715	0.000	
	S3	45.2 ± 9.3	(21.5 - 75.1)	40.4 ± 8.6	(21.5 - 61.1)	45.7 ± 8.4	(28.4 - 69.2)	49.5 ± 8.9	(31.3 - 75.1)	32.509	0.000	
	S2	32.0 ± 7.1	(11.9 - 53.6)	31.4 ± 7.1	(12.2 - 49.0)	32.1 ± 6.5	(13.4 - 53.6)	32.5 ± 7.9	(11.9 - 52.9)	0.626	0.535	
	S1	27.9 ± 9.9	(-0.7 - 61.8)	34.2 ± 9.9	(14.0 - 61.8)	28.2 ± 8.3	(8.2 - 52.2)	21.0 ± 7.3	(-0.7 - 35.6)	68.121	0.000	
	L5	74.3 ± 8.5	(51.0 - 96.0)	80.6 ± 6.8	(63.6 - 96.0)	74.2 ± 6.9	(53.6 - 90.0)	67.8 ± 7.2	(51.0 - 88.9)	98.166	0.000	
	L4	91.0 ± 7.4	(68.4 - 111.2)	95.2 ± 6.6	(83.4 - 111.2)	90.6 ± 6.7	(68.4 - 105.1)	87.1 ± 6.9	(69.7 - 107.1)	41.902	0.000	
	L3	100.9 ± 6.9	(79.3 - 122.6)	103.8 ± 7.0	(87.0 - 122.6)	100.3 ± 6.6	(79.3 - 113.9)	98.8 ± 6.3	(84.6 - 117.3)	17.609	0.000	
	L2	108.0 ± 6.6	(88.5 - 131.0)	109.2 ± 6.8	(95.5 - 131.0)	107.9 ± 6.6	(88.5 - 124.8)	107.0 ± 6.1	(92.1 - 121.4)	3.397	0.034	
	L1	108.6 ± 6.3	(84.8 - 130.9)	108.4 ± 6.9	(84.8 - 126.6)	108.5 ± 6.2	(91.0 - 130.9)	109.0 ± 5.9	(96.1 - 125.8)	0.331	0.718	
	T12	107.8 ± 5.9	(88.0 - 126.9)	106.9 ± 6.3	(88.0 - 125.5)	107.7 ± 5.8	(90.0 - 126.9)	108.8 ± 5.4	(96.6 - 121.7)	2.955	0.053	
	T11	105.1 ± 5.8	(89.4 - 121.8)	103.8 ± 5.9	(91.7 - 120.3)	105.1 ± 5.6	(89.4 - 120.1)	106.5 ± 5.8	(91.9 - 121.8)	6.248	0.002	
	T10	102.8 ± 5.6	(86.6 - 121.2)	101.8 ± 5.5	(90.0 - 117.5)	102.8 ± 5.9	(86.6 - 121.2)	104.0 ± 5.1	(88.0 - 119.5)	4.847	0.008	
	T9	101.7 ± 5.6	(84.9 - 119.0)	100.7 ± 5.5	(90.0 - 117.9)	101.8 ± 5.7	(84.9 - 116.7)	102.7 ± 5.5	(85.6 - 119.0)	3.966	0.020	
	T8	99.6 ± 6.0	(82.5 - 116.6)	98.8 ± 6.2	(85.2 - 116.6)	99.6 ± 5.9	(82.5 - 112.0)	100.4 ± 5.8	(87.1 - 113.5)	2.130	0.120	
	T7	95.9 ± 5.8	(79.1 - 112.3)	94.5 ± 5.7	(81.6 - 108.9)	96.4 ± 5.9	(81.9 - 112.3)	96.7 ± 5.8	(79.1 - 111.5)	5.057	0.007	
	T6	90.7 ± 6.0	(72.7 - 109.3)	89.7 ± 6.0	(76.9 - 102.0)	90.9 ± 6.0	(73.2 - 106.0)	91.3 ± 6.1	(72.7 - 109.3)	2.393	0.093	
	T5	84.9 ± 6.7	(67.9 - 104.5)	83.7 ± 6.7	(67.9 - 96.9)	85.2 ± 6.3	(69.5 - 99.8)	85.8 ± 7.0	(68.0 - 104.5)	3.097	0.046	
	T4	80.1 ± 7.2	(52.8 - 103.4)	79.5 ± 7.1	(60.0 - 96.3)	80.1 ± 7.0	(55.5 - 98.0)	80.6 ± 7.7	(52.8 - 103.4)	0.670	0.512	
	T3	75.9 ± 7.5	(53.3 - 101.7)	75.5 ± 7.7	(53.5 - 97.9)	75.9 ± 7.3	(53.3 - 101.7)	76.4 ± 7.6	(53.5 - 95.7)	0.426	0.654	
	T2	72.5 ± 7.9	(51.8 - 101.2)	72.4 ± 7.8	(52.6 - 93.6)	72.0 ± 8.0	(52.1 - 101.2)	73.4 ± 8.0	(51.8 - 90.0)	0.965	0.382	
	T1	71.5 ± 8.4	(48.9 - 94.3)	71.3 ± 8.4	(48.9 - 93.5)	70.9 ± 8.4	(52.2 - 93.4)	72.4 ± 8.4	(53.9 - 94.3)	1.114	0.329	
	C7	79.0 ± 8.6	(54.7 - 108.6)	77.9 ± 8.7	(54.7 - 101.6)	79.4 ± 8.6	(56.8 - 108.6)	79.5 ± 8.6	(56.6 - 103.3)	1.394	0.249	
	Ferguson LI-S2	76.6 ± 9.8	(39.1 - 109.7)	76.9 ± 10.4	(50.3 - 109.7)	76.4 ± 9.6	(39.1 - 100.5)	76.5 ± 9.7	(55.5 - 101.6)	0.099	0.906	
	Ferguson T1-T12	36.3 ± 10.9	(5.1 - 71.8)	35.6 ± 11.7	(6.9 - 71.8)	36.8 ± 10.7	(5.1 - 67.0)	36.3 ± 10.5	(7.8 - 63.7)	0.416	0.660	
	Smin	25.5 ± 7.5	(-0.7 - 44.5)	28.8 ± 6.7	(12.2 - 44.5)	26.4 ± 6.7	(8.2 - 47.0)	20.8 ± 7.1	(-0.7 - 35.5)	41.771	0.000	
	Tmax	111.7 ± 5.4	(96.3 - 131.0)	111.9 ± 5.9	(101.4 - 131.0)	111.4 ± 5.2	(96.3 - 130.9)	111.7 ± 5.0	(102.1 - 125.8)	0.316	0.729	
	Tmin	70.0 ± 7.8	(48.9 - 92.3)	69.7 ± 7.8	(48.9 - 93.2)	69.5 ± 7.8	(52.1 - 92.3)	70.8 ± 7.8	(51.8 - 90.0)	1.087	0.338	
	Ferguson L5Lmax	86.2 ± 9.8	(58.7 - 118.7)	83.2 ± 10.0	(62.8 - 118.7)	85.0 ± 9.0	(58.7 - 109.3)	90.9 ± 9.0	(70.7 - 110.3)	21.899	0.000	
	Ferguson T1Kmax	41.7 ± 10.3	(12.6 - 77.2)	42.2 ± 11.1	(16.7 - 77.2)	42.0 ± 9.9	(17.2 - 67.0)	40.9 ± 9.9	(12.6 - 71.2)	0.558	0.573	
	middle of the vertebral body	Co1	96.9 ± 13.4	(54.4 - 139.6)	90.4 ± 12.2	(58.6 - 121.7)	97.1 ± 12.0	(54.4 - 135.0)	103.3 ± 13.5	(72.1 - 139.6)	30.818	0.000
S5		81.4 ± 11.8	(44.0 - 113.1)	73.6 ± 11.4	(44.0 - 97.9)	82.2 ± 9.8	(55.2 - 103.7)	88.2 ± 9.9	(59.2 - 113.1)	57.758	0.000	
S4		64.2 ± 12.0	(30.1 - 92.5)	56.5 ± 10.3	(33.5 - 86.7)	64.2 ± 10.5	(30.1 - 84.3)	72.1 ± 10.6	(47.3 - 92.5)	64.047	0.000	
S3		49.7 ± 9.1	(27.2 - 78.0)	44.5 ± 8.3	(27.2 - 63.9)	50.0 ± 8.0	(32.4 - 71.9)	54.5 ± 8.5	(34.2 - 78.0)	42.145	0.000	
S2		37.5 ± 7.0	(15.1 - 58.0)	36.8 ± 7.0	(15.1 - 53.3)	37.8 ± 6.5	(19.0 - 55.9)	37.8 ± 7.7	(16.2 - 58.0)	0.875	0.418	
S1		37.0 ± 9.1	(8.2 - 67.8)	42.3 ± 9.1	(19.4 - 67.8)	37.3 ± 7.6	(17.6 - 62.0)	31.0 ± 7.2	(8.2 - 47.0)	57.806	0.000	
L5		75.8 ± 8.6	(39.6 - 98.2)	81.8 ± 7.7	(39.6 - 98.2)	75.8 ± 6.9	(56.0 - 92.7)	69.5 ± 7.2	(52.8 - 88.7)	82.188	0.000	
L4		91.6 ± 7.4	(70.1 - 111.5)	95.9 ± 6.7	(81.6 - 111.5)	91.1 ± 6.6	(70.1 - 106.1)	87.8 ± 6.8	(71.7 - 105.4)	42.939	0.000	
L3		101.3 ± 7.0	(78.0 - 123.4)	104.1 ± 7.1	(85.6 - 123.4)	100.7 ± 6.6	(78.0 - 115.0)	99.2 ± 6.4	(83.6 - 119.6)	16.270	0.000	
L2		107.2 ± 6.5	(87.6 - 132.0)	108.2 ± 6.8	(93.9 - 132.0)	107.0 ± 6.5	(87.6 - 126.0)	106.4 ± 6.1	(90.7 - 121.5)	2.254	0.106	
L1		107.8 ± 6.3	(85.1 - 129.6)	107.6 ± 6.8	(85.1 - 127.1)	107.6 ± 6.1	(90.4 - 129.6)	108.2 ± 5.9	(96.5 - 124.2)	0.402	0.669	
T12		106.5 ± 6.0	(85.0 - 125.8)	105.4 ± 6.4	(85.0 - 125.8)	106.2 ± 5.9	(88.7 - 125.5)	107.9 ± 5.5	(96.3 - 122.4)	5.215	0.006	
T11		104.1 ± 5.8	(88.5 - 122.5)	102.7 ± 5.8	(89.3 - 120.8)	104.0 ± 5.7	(88.5 - 117.9)	105.7 ± 5.6	(91.4 - 122.5)	8.154	0.000	
T10		101.9 ± 5.6	(85.3 - 119.4)	100.9 ± 5.5	(88.9 - 119.4)	101.9 ± 5.7	(85.3 - 117.5)	103.1 ± 5.1	(89.1 - 118.1)	4.905	0.008	
T9		100.9 ± 5.7	(84.2 - 117.6)	99.8 ± 5.7	(87.8 - 117.6)	100.9 ± 5.8	(84.2 - 117.2)	102.0 ± 5.4	(84.2 - 115.3)	4.103	0.017	
T8		98.6 ± 5.8	(82.9 - 115.4)	97.7 ± 6.0	(85.2 - 115.4)	98.6 ± 5.7	(82.9 - 111.9)	99.4 ± 5.6	(85.5 - 112.0)	2.606	0.075	
T7		94.4 ± 5.8	(78.5 - 109.6)	93.0 ± 5.5	(79.4 - 106.5)	94.8 ± 5.8	(80.3 - 109.0)	95.4 ± 5.8	(78.5 - 109.6)	5.590	0.004	
T6		88.9 ± 5.8	(72.2 - 108.2)	87.9 ± 5.8	(75.9 - 108.8)	89.2 ± 5.8	(72.4 - 101.1)	89.4 ± 5.8	(72.2 - 108.2)	2.489	0.084	
T5		83.1 ± 6.6	(64.8 - 102.5)	82.0 ± 6.6	(67.3 - 95.9)	83.3 ± 6.2	(69.2 - 96.3)	83.9 ± 6.9	(64.8 - 102.5)	2.629	0.073	
T4		78.1 ± 7.1	(53.0 - 100.5)	77.6 ± 6.9	(61.5 - 93.6)	78.1 ± 7.0	(55.8 - 99.2)	78.5 ± 7.4	(53.0 - 100.5)	0.554	0.575	
T3		73.5 ± 7.5	(50.3 - 99.1)	73.2 ± 7.8	(53.1 - 95.1)	73.3 ± 7.1	(50.3 - 99.1)	74.1 ± 7.7	(53.4 - 92.8)	0.569	0.566	
T2		70.3 ± 8.0	(47.2 - 94.2)	70.1 ± 8.0	(47.2 - 91.8)	69.8 ± 8.0	(49.2 - 94.2)	71.2 ± 7.9	(51.3 - 89.3)	0.984	0.375	
T1		69.6 ± 8.6	(46.0 - 93.0)	69.4 ± 8.5	(46.0 - 93.0)	69.2 ± 8.6	(48.2 - 90.0)	70.3 ± 8.8	(46.9 - 91.6)	0.661	0.517	
C7		78.8 ± 8.5	(52.2 - 109.1)	77.8 ± 8.7	(52.2 - 102.3)	79.1 ± 8.6	(56.3 - 109.1)	79.3 ± 8.2	(56.5 - 101.3)	1.143	0.320	
Ferguson LI-S2		70.3 ± 9.9	(37.0 - 106.2)	70.8 ± 10.4	(46.9 - 106.2)	69.8 ± 9.7	(37.0 - 95.4)	70.4 ± 9.6	(49.3 - 95.8)	0.391	0.677	
Ferguson T1-T12		36.9 ± 11.1	(5.0 - 75.4)	36.0 ± 11.9	(5.0 - 75.4)	37.0 ± 10.9	(5.5 - 66.0)	37.6 ± 10.7	(7.4 - 66.3)	0.569	0.567	
Smin		33.8 ± 7.0	(8.2 - 52.4)	35.5 ± 6.9	(15.1 - 52.4)	34.9 ± 6.3	(17.6 - 48.9)	30.5 ± 7.0	(8.2 - 46.7)	19.839	0.000	
Tmax		110.5 ± 5.5	(94.4 - 132.0)	110.8 ± 6.0	(100.6 - 132.0)	110.2 ± 5.3	(94.4 - 129.6)	110.6 ± 5.2	(99.4 - 124.2)	0.443	0.642	
Tmin		68.1 ± 8.0	(46.0 - 89.4)	67.9 ± 8.1	(46.0 - 89.4)	67.7 ± 8.0	(48.2 - 88.6)	68.9 ± 8.0	(46.9 - 88.6)	0.799	0.450	
Ferguson L5Lmax		76.7 ± 9.6	(45.5 - 116.9)	75.3 ± 10.3	(53.9 - 116.9)	75.2 ± 8.9	(45.5 - 98.2)	80.1 ± 8.8	(57.7 - 101.1)	10.674	0.000	
Ferguson T1Kmax		42.3 ± 10.6	(14.0 - 80.7)	42.9 ± 11.5	(16.7 - 80.7)	42.4 ± 10.3	(17.2 - 70.0)	41.7 ± 10.1	(14.0 - 70.5)	0.394	0.674	
posterior edge of the vertebral body		Co1	110.8 ± 14.1	(65.6 - 164.7)	104.0 ± 12.9	(75.3 - 140.7)	111.0 ± 12.5	(65.6 - 150.5)	117.5 ± 14.2	(87.5 - 164.7)	30.134	0.000
		S5	82.2 ± 12.0	(39.5 - 110.9)	74.3 ± 11.5	(39.5 - 96.7)	83.0 ± 10.2	(56.0 - 105.6)	89.1 ± 10.1	(59.3 - 110.9)	56.909	0.000
	S4	66.6 ± 11.6	(34.5 - 96.1)	59.4 ± 10.0	(34.8 - 88.9)	66.7 ± 10.1	(34.5 - 87.1)	73.8 ± 10.5	(49.0 - 96.1)	57.090	0.000	
	S3	53.7 ± 9.3	(28.6 - 80.1)	48.5 ± 8.4	(28.6 - 67.8)	53.9 ± 8.1	(35.4 - 76.8)	58.9 ± 8.7	(36.7 - 80.1)	44.390	0.000	
	S2	43.1 ± 7.5	(18.1 - 64.2)	42.5 ± 7.5	(18.1 - 61.2)	43.7 ± 7.2	(22.6 - 64.2)	42.9 ± 7.9	(20.3 - 62.6)	0.863	0.423	
	S1	48.2 ± 8.7	(17.5 - 75.1)	52.7 ± 8.9								

Table 3 Classification based on TLmax - vertebral tilt and Ferguson angle of sagittal spinal alignment

	Total (n=388)			Classification of TLmax						F	P
				Group A (n=91)		Group A (n=216)		Group C (n=81)			
TLmax	110.5 ± 5.5	(94.4 - 132.0)		104.0 ± 2.3	(94.4 - 106.5)	110.2 ± 2.1	(106.6 - 114.5)	118.6 ± 3.8	(114.7 - 132.0)	700.322	0.000
anterior edge of the vertebral body											
Co1	82.0 ± 13.5	(43.2 - 117.7)		80.3 ± 14.1	(47.6 - 117.7)	83.0 ± 13.8	(43.2 - 117.6)	81.2 ± 11.8	(52.9 - 109.9)	1.434	0.240
S5	80.4 ± 12.0	(40.7 - 117.7)		80.3 ± 12.2	(40.7 - 108.4)	80.7 ± 12.1	(50.2 - 117.7)	80.0 ± 11.7	(45.5 - 108.4)	0.091	0.913
S4	61.6 ± 12.9	(25.7 - 92.9)		61.5 ± 13.2	(27.0 - 89.1)	62.3 ± 12.5	(31.1 - 92.9)	59.8 ± 13.7	(25.7 - 86.6)	1.041	0.354
S3	45.2 ± 9.3	(21.5 - 75.1)		44.5 ± 9.3	(23.4 - 66.7)	45.8 ± 9.4	(21.5 - 69.2)	44.5 ± 9.2	(25.6 - 75.1)	0.860	0.424
S2	32.0 ± 7.1	(11.9 - 53.6)		32.0 ± 7.4	(13.4 - 53.6)	32.4 ± 7.1	(11.9 - 52.9)	30.9 ± 6.6	(12.2 - 47.4)	1.383	0.252
S1	27.9 ± 9.9	(-0.7 - 61.8)		30.3 ± 11.5	(-0.7 - 60.1)	27.7 ± 9.7	(5.5 - 61.8)	25.8 ± 8.1	(8.2 - 47.8)	4.590	0.011
L5	74.3 ± 8.5	(51.0 - 96.0)		74.4 ± 9.1	(52.6 - 96.0)	73.2 ± 8.4	(51.0 - 93.2)	77.0 ± 7.8	(57.6 - 95.7)	5.805	0.003
L4	91.0 ± 7.4	(68.4 - 111.2)		88.7 ± 7.3	(68.4 - 102.9)	90.1 ± 6.7	(73.5 - 108.4)	96.0 ± 7.2	(69.7 - 111.2)	27.928	0.000
L3	100.9 ± 6.9	(79.3 - 122.6)		96.6 ± 5.5	(79.3 - 106.6)	100.5 ± 5.9	(84.6 - 112.9)	106.9 ± 6.8	(91.9 - 122.6)	65.479	0.000
L2	108.0 ± 6.6	(88.5 - 131.0)		102.1 ± 4.4	(88.5 - 110.2)	107.7 ± 4.5	(92.1 - 117.6)	115.5 ± 6.0	(95.5 - 131.0)	168.117	0.000
L1	108.6 ± 6.3	(84.8 - 130.9)		102.1 ± 4.0	(88.0 - 107.7)	108.5 ± 3.7	(97.0 - 118.6)	116.2 ± 5.6	(92.5 - 130.9)	240.672	0.000
T12	107.8 ± 5.9	(88.0 - 126.9)		102.0 ± 4.0	(88.0 - 107.9)	108.0 ± 4.0	(95.4 - 116.1)	113.6 ± 5.9	(97.1 - 126.9)	146.757	0.000
T11	105.1 ± 5.8	(89.4 - 121.8)		100.8 ± 4.1	(89.4 - 107.7)	105.2 ± 4.7	(91.9 - 115.4)	109.7 ± 6.6	(93.5 - 121.8)	67.544	0.000
T10	102.8 ± 5.6	(86.6 - 121.2)		99.7 ± 4.5	(87.1 - 108.4)	103.3 ± 5.0	(86.6 - 114.6)	105.2 ± 6.6	(91.6 - 121.2)	25.202	0.000
T9	101.7 ± 5.6	(84.9 - 119.0)		99.7 ± 4.4	(84.9 - 108.4)	102.2 ± 5.4	(85.6 - 114.3)	102.8 ± 6.8	(89.1 - 119.0)	8.767	0.000
T8	99.6 ± 6.0	(82.5 - 116.6)		98.4 ± 4.5	(82.5 - 108.6)	100.0 ± 5.9	(86.1 - 113.5)	99.7 ± 7.4	(85.2 - 116.6)	2.337	0.098
T7	95.9 ± 5.8	(79.1 - 112.3)		95.3 ± 4.6	(82.2 - 106.4)	96.6 ± 5.8	(79.1 - 112.3)	94.8 ± 6.9	(81.6 - 109.0)	3.112	0.046
T6	90.7 ± 6.0	(72.7 - 109.3)		90.8 ± 5.0	(78.5 - 103.7)	91.1 ± 6.3	(72.7 - 109.3)	89.2 ± 6.2	(77.5 - 101.4)	3.016	0.050
T5	84.9 ± 6.7	(67.9 - 104.5)		85.6 ± 5.9	(70.9 - 98.7)	85.3 ± 6.7	(68.0 - 104.5)	83.3 ± 7.2	(67.9 - 99.8)	3.291	0.038
T4	80.1 ± 7.2	(52.8 - 103.4)		81.6 ± 6.9	(64.6 - 98.0)	80.3 ± 7.1	(52.8 - 103.4)	77.9 ± 7.6	(60.0 - 94.0)	5.727	0.004
T3	75.9 ± 7.5	(53.3 - 101.7)		77.8 ± 7.3	(61.2 - 101.7)	75.9 ± 7.4	(53.3 - 97.9)	73.8 ± 7.6	(53.5 - 89.1)	6.302	0.002
T2	72.5 ± 7.9	(51.8 - 101.2)		74.3 ± 8.0	(51.8 - 101.2)	72.6 ± 7.8	(52.1 - 93.6)	70.3 ± 7.8	(52.9 - 87.9)	5.551	0.004
T1	71.5 ± 8.4	(48.9 - 94.3)		73.3 ± 7.9	(55.3 - 94.3)	71.1 ± 8.4	(49.7 - 93.5)	70.2 ± 8.7	(48.9 - 90.6)	3.341	0.036
C7	79.0 ± 8.6	(54.7 - 108.6)		80.1 ± 8.3	(61.5 - 103.3)	78.6 ± 8.4	(56.6 - 108.6)	78.8 ± 9.5	(54.7 - 99.7)	0.995	0.371
Ferguson L1-S2	76.6 ± 9.8	(39.1 - 109.7)		70.1 ± 8.8	(39.1 - 88.7)	76.1 ± 8.1	(55.8 - 97.1)	85.4 ± 8.8	(61.8 - 109.7)	70.926	0.000
Ferguson T1-T12	36.3 ± 10.9	(5.1 - 71.8)		28.7 ± 9.7	(5.1 - 50.0)	36.8 ± 9.4	(14.7 - 60.3)	43.4 ± 10.7	(11.3 - 71.8)	49.679	0.000
Smin	25.5 ± 7.5	(-0.7 - 44.5)		26.5 ± 8.4	(-0.7 - 44.0)	25.6 ± 7.5	(5.5 - 44.5)	23.9 ± 6.2	(8.2 - 37.5)	2.653	0.072
TLmax	111.7 ± 5.4	(96.3 - 131.0)		105.4 ± 2.4	(96.3 - 110.2)	111.4 ± 2.3	(104.0 - 118.6)	119.4 ± 3.6	(114.0 - 131.0)	598.424	0.000
Tmin	70.0 ± 7.8	(48.9 - 92.3)		71.7 ± 7.5	(51.8 - 92.3)	69.9 ± 7.8	(49.7 - 91.2)	68.3 ± 7.9	(48.9 - 86.7)	4.267	0.015
Ferguson L5Lmax	86.2 ± 9.8	(58.7 - 118.7)		78.9 ± 9.4	(58.7 - 108.6)	85.8 ± 8.0	(65.2 - 108.2)	95.5 ± 7.0	(79.9 - 118.7)	89.283	0.000
Ferguson TKmax	41.7 ± 10.3	(12.6 - 77.2)		33.7 ± 7.7	(12.6 - 54.4)	41.6 ± 8.4	(20.9 - 63.9)	51.1 ± 9.5	(29.8 - 77.2)	90.206	0.000
middle of the vertebral body											
Co1	96.9 ± 13.4	(54.4 - 139.6)		95.6 ± 13.4	(62.6 - 135.0)	97.7 ± 13.9	(54.4 - 139.6)	96.1 ± 12.0	(70.6 - 124.4)	0.906	0.405
S5	81.4 ± 11.8	(44.0 - 113.1)		80.9 ± 12.2	(44.0 - 103.9)	81.8 ± 11.7	(53.3 - 113.1)	80.5 ± 11.7	(45.7 - 104.8)	0.444	0.642
S4	64.2 ± 12.0	(30.1 - 92.5)		64.0 ± 11.8	(36.1 - 89.5)	64.7 ± 11.8	(33.5 - 92.5)	63.0 ± 12.9	(30.1 - 89.6)	0.642	0.527
S3	49.7 ± 9.1	(27.2 - 78.0)		49.1 ± 9.3	(28.3 - 71.9)	50.2 ± 9.0	(27.3 - 75.1)	48.8 ± 9.1	(27.2 - 78.0)	0.955	0.386
S2	37.5 ± 7.0	(15.1 - 58.0)		37.8 ± 7.2	(21.0 - 55.9)	38.0 ± 7.0	(16.2 - 58.0)	35.9 ± 6.7	(15.1 - 50.7)	2.929	0.055
S1	37.0 ± 9.1	(8.2 - 67.8)		39.0 ± 10.3	(8.2 - 65.7)	36.9 ± 8.9	(16.1 - 67.8)	34.8 ± 7.6	(17.2 - 54.9)	4.556	0.011
L5	75.8 ± 8.6	(39.6 - 98.2)		75.8 ± 8.9	(52.8 - 95.9)	74.6 ± 8.6	(39.6 - 94.5)	78.8 ± 7.6	(61.0 - 98.2)	7.165	0.001
L4	91.6 ± 7.4	(70.1 - 111.5)		89.2 ± 7.2	(70.1 - 103.2)	90.8 ± 6.6	(73.7 - 108.4)	96.6 ± 7.3	(71.7 - 111.5)	28.669	0.000
L3	101.3 ± 7.0	(80.8 - 123.4)		96.8 ± 5.6	(78.0 - 106.2)	100.8 ± 5.8	(83.5 - 112.9)	107.7 ± 6.8	(92.6 - 123.4)	74.191	0.000
L2	107.2 ± 6.5	(87.6 - 132.0)		100.8 ± 4.0	(87.6 - 106.4)	107.0 ± 4.0	(90.7 - 114.3)	115.0 ± 5.8	(93.9 - 132.0)	219.535	0.000
L1	107.8 ± 6.3	(85.1 - 129.6)		101.4 ± 3.9	(85.1 - 106.5)	107.6 ± 3.5	(96.0 - 114.5)	115.5 ± 5.8	(93.3 - 129.6)	250.312	0.000
T12	106.5 ± 6.0	(85.0 - 125.8)		100.7 ± 4.1	(85.0 - 106.0)	106.6 ± 4.0	(93.7 - 114.4)	112.5 ± 6.2	(95.5 - 125.8)	143.972	0.000
T11	104.1 ± 5.8	(88.5 - 122.5)		99.7 ± 4.1	(88.5 - 106.5)	104.4 ± 4.7	(90.3 - 113.4)	108.4 ± 6.6	(92.2 - 122.5)	64.953	0.000
T10	101.9 ± 5.6	(85.3 - 119.4)		98.8 ± 4.4	(85.3 - 106.3)	102.4 ± 5.0	(85.5 - 113.6)	104.3 ± 6.5	(91.7 - 119.4)	25.332	0.000
T9	100.9 ± 5.7	(84.2 - 117.6)		98.9 ± 4.1	(84.2 - 106.5)	101.4 ± 5.6	(84.2 - 114.1)	101.8 ± 7.0	(87.5 - 117.6)	7.812	0.000
T8	98.6 ± 5.8	(82.9 - 115.4)		97.4 ± 4.3	(82.9 - 106.5)	99.1 ± 5.7	(85.3 - 112.0)	98.7 ± 7.3	(84.2 - 115.4)	2.801	0.062
T7	94.4 ± 5.8	(78.5 - 109.6)		93.8 ± 4.5	(81.4 - 104.0)	95.1 ± 5.7	(78.5 - 109.6)	93.3 ± 6.9	(79.4 - 108.3)	3.431	0.033
T6	88.9 ± 5.8	(72.2 - 108.2)		88.8 ± 5.0	(76.5 - 98.5)	89.5 ± 6.0	(72.2 - 108.2)	87.4 ± 6.1	(75.9 - 98.9)	3.854	0.022
T5	83.1 ± 6.6	(64.8 - 102.5)		83.9 ± 6.1	(66.0 - 97.9)	83.5 ± 6.5	(64.8 - 102.5)	81.1 ± 7.0	(67.3 - 96.1)	4.855	0.008
T4	78.1 ± 7.1	(53.0 - 100.5)		79.7 ± 6.9	(64.7 - 99.2)	78.2 ± 7.0	(53.0 - 100.5)	75.9 ± 7.0	(58.8 - 90.0)	6.567	0.002
T3	73.5 ± 7.5	(50.3 - 99.1)		75.6 ± 7.4	(57.8 - 99.1)	73.4 ± 7.3	(50.3 - 95.1)	71.3 ± 7.5	(53.1 - 85.9)	7.308	0.001
T2	70.3 ± 8.0	(47.2 - 94.2)		72.2 ± 7.8	(51.3 - 94.2)	70.2 ± 7.9	(47.2 - 91.8)	68.3 ± 8.0	(50.2 - 85.9)	5.209	0.006
T1	69.6 ± 8.6	(46.0 - 93.0)		71.6 ± 8.0	(53.7 - 91.6)	69.2 ± 8.6	(46.9 - 93.0)	68.4 ± 9.0	(46.0 - 89.1)	3.466	0.032
C7	78.8 ± 8.5	(52.2 - 109.1)		79.9 ± 8.1	(63.4 - 101.3)	78.5 ± 8.4	(56.3 - 109.1)	78.2 ± 9.4	(52.2 - 97.2)	1.157	0.315
Ferguson L1-S2	70.3 ± 9.9	(37.0 - 106.2)		63.6 ± 8.4	(37.0 - 82.0)	69.6 ± 8.0	(49.3 - 89.4)	79.7 ± 9.1	(58.6 - 106.2)	81.959	0.000
Ferguson T1-T12	36.9 ± 11.1	(5.0 - 75.4)		29.2 ± 10.1	(5.0 - 50.9)	37.4 ± 9.4	(15.2 - 61.3)	44.2 ± 11.3	(11.0 - 75.4)	48.970	0.000
Smin	33.8 ± 7.0	(8.2 - 52.4)		34.7 ± 7.9	(8.2 - 52.0)	34.1 ± 6.8	(16.1 - 52.4)	32.0 ± 6.0	(15.1 - 46.7)	3.764	0.024
TLmax	110.5 ± 5.5	(94.4 - 132.0)		104.0 ± 2.3	(94.4 - 106.5)	110.2 ± 2.1	(106.6 - 114.5)	118.6 ± 3.8	(114.7 - 132.0)	700.322	0.000
Tmin	68.1 ± 8.0	(46.0 - 89.4)		70.1 ± 7.6	(51.3 - 86.6)	68.0 ± 8.0	(46.9 - 89.4)	66.4 ± 8.2	(46.0 - 84.6)	4.894	0.008
Ferguson L5Lmax	76.7 ± 9.6	(45.5 - 116.9)		69.4 ± 8.7	(45.5 - 97.8)	76.0 ± 7.3	(59.1 - 93.9)	86.7 ± 7.3	(69.9 - 116.9)	111.514	0.000
Ferguson TKmax	42.3 ± 10.6	(14.0 - 80.7)		33.9 ± 8.1	(14.0 - 54.3)	42.2 ± 8.4	(21.3 - 65.6)	52.2 ± 9.9	(32.7 - 80.7)	95.900	0.000
posterior edge of the vertebral body											
Co1	110.8 ± 14.1	(65.6 - 164.7)		110.0 ± 13.5	(75.3 - 150.5)	111.4 ± 14.6	(65.6 - 164.7)	110.1 ± 13.6	(82.3 - 150.1)	0.419	0.658
S5	82.2 ± 12.0	(39.5 - 110.9)		81.5 ± 12.6	(39.5 - 105.6)	82.9 ± 11.7	(54.5 - 110.9)	81.0 ± 12.1	(45.9 - 108.4)	0.934	0.394
S4	66.6 ± 11.6	(34.5 - 96.1)		66.2 ± 10.9	(41.4 - 93.0)	67.0 ± 11.6	(34.8 - 93.2)	65.8 ± 12.5	(34		

1. Classification based on Pelvic Incidence (PI)

According to the anatomical segmentation, the study results show that for groups with different types of PI, there are no statistically significant differences in the vertebral tilt of the transitional vertebrae S2, L1, T1 for the "S" curve anatomical segmentation (Groups A, B, C), nor are there differences in the lumbosacral lordosis (LSL) Ferguson L1-S2 and thoracic kyphosis (TK) Ferguson T1-T12. These results imply that PI does not significantly affect the overall magnitude of the lumbosacral lordosis and thoracic kyphosis; in other words, the overall curvature of the "S" curve of the spine is essentially unaffected by PI according to anatomical segmentation. (Table 1, Figure 1)

Based on functional segmentation, there are no significant statistical differences in TLmax and Tmin; Ferguson TKmax also shows no significant statistical difference. There is a statistical difference in the Smin and Ferguson LSLmax. The study results indicate that it is more common for the anterior edge of Smin to be located at S1, while the posterior edge of Smin is mainly concentrated at S2, with the middle part of the vertebrae falling between the two. Unlike the thoracolumbar vertebrae, which are rectangular in shape, the sacral vertebrae are trapezoidal, with S1 having a greater trapezoidal gradient than S2. The variation in the anterior edge tilt is greater than that of the posterior edge, which could significantly affect the difference between measurement results and the true sequence. As the results show, the anterior edge of Smin has a greater variation than the middle and posterior edges, and although there is a statistical difference in the posterior edge ($0.01 < P < 0.05$), the significance is lower than that of the anterior and middle edges ($P < 0.001$), and the mean difference is small, even within the acceptable measurement error range. Based on these inferences, the author believes that due to the specific morphology of S1, even with the use of vertebral tilt measurements and the Ferguson method, the measurement results may not accurately reflect its sagittal alignment, and this could also likely lead to the statistical differences in Smin and Ferguson LSLmax observed in this study. Although not proven, based on the logical analysis of the results, the author leans towards the belief that the true Smin and Ferguson LSLmax also do not show statistical differences. If that were the case, according to functional segmentation, the overall curvature of the spinal "S" curve is also unaffected by PI. (Table 1, Figure 2)

Figure 1 Classification based on PI - anatomical segmentation





Figure 2 Classification based on PI - functional segmentation



The study results also show that for groups with different types of PI, there are varying degrees of statistical differences or trends in S1, L5, L4, L3, L2, T12, T11, T10, T9, T7, with more significant differences in the lumbosacral segment. This means that although the overall curvature of the "S" curve of the spine is the same across different PI groups, there are differences in local curvature. The author will discuss the mechanisms and logic behind this in subsequent articles.

2. Classification based on the maximal thoracolumbar vertebral tilt (TLmax)

Regardless of whether anatomical or functional segmentation is considered, this study shows that for groups with different TLmax, there are significant statistical differences in the tilt of the transitional vertebrae S2, L1, T1 for the "S" curve anatomical segmentation, as well as the transitional vertebrae Smin, TLmax, Tmin for functional segmentation (Groups A, B, C). There are also significant statistical differences in LSL, (Ferguson L1-S2), TK (Ferguson T1-T12), Ferguson LSLmax, and Ferguson TKmax. These results imply that using TLmax as a classification reference can effectively differentiate the overall curvature of the "S" curve of the human spine. (Table 3, Figure 3,4)

Figure 3 Classification based on TLmax - functional segmentation





Figure 4 Classification based on TLmax - anatomical segmentation



At the same time, S3 and C7, as adjacent segments of the "S" curve, show no significant statistical differences in their tilts. This result means that the curvature magnitude of the "S" curve does not have a significant impact on the regional range of curve. Additionally, there are no significant statistical differences in PI and SS across the groups of this classification method, which further proves that PI, and even SS, do not substantially decide the overall curvature magnitude of the spinal "S" curve.

The classification method using TLmax as a reference provides an alternative perspective for our research on the sagittal balance of the human spine, which could potentially influence the diagnosis and treatment strategies for disorders of the spinal sagittal alignment.

Limitations of the study: 1. The study utilized a multicenter sample, and for a subset of participants, data on gender and precise age were absent. 2. PI is a morphological parameter. The author had intended to contrast and categorize using newly developed morphological parameters, but the description and interpretation of the parameter would necessitate a substantial exposition. To ensure a straightforward and coherent narrative, this paper provisionally employs the positional parameter TLmax as an interim classification method to avoid reader confusion. 3. The sample sizes for the two categorical groups in this study were

not uniform. This is because uniform sample sizes across all categories and groups did not yield sufficiently significant statistical outcomes, which could potentially perplex readers. The study's approach to sample categorization aims to clarify results but may inherently introduce a degree of logical stringency imperfection.

Conclusion

Pelvic Incidence (PI) does not influence the overall magnitude of lumbosacral lordosis (LSL) and thoracic kyphosis (TK). The sagittal spinal classification based on PI reveals no difference in the overall curvature of the spinal "S" shape. The maximum thoracic kyphosis (TLmax) significantly affects LSL and TK, and it can be considered a straightforward radiological parameter for distinguishing the overall curvature of the spinal "S" shape.

References

1. Le Huec, J. C., Faundez, A., Dominguez, D., et al. (2015). Evidence showing the relationship between sagittal balance and clinical outcomes in surgical treatment of degenerative spinal diseases: a literature review. *International Orthopaedics*, 39, 87-95. <https://doi.org/10.1007/s00264-014-2516-6>
2. Diebo, B. G., Varghese, J. J., Lafage, R., et al. (2015). Sagittal alignment of the spine: What do you need to know? *Clinical Neurology and Neurosurgery*, 139, 139. <https://doi.org/10.1016/j.clineuro.2015.10.024>
3. Lovejoy, C. O. (2005). The natural history of human gait and posture. Part 1. Spine and pelvis. *Gait & Posture*, 21(1), 95-112.
4. Farfan, H. F. (1978). The biomechanical advantage of lordosis and hip extension for upright activity. Man as compared with other anthropoids. *Spine*, 3(4), 336-342.
5. BEEN, E., GOMEZ-OLIVENCIA, A., SHEFI, S., et al. (2017). Evolution of Spinopelvic Alignment in Hominins. *The Anatomical Record: Advances in Integrative Anatomy and Evolutionary Biology*, 300(5), 900-911. <https://doi.org/10.1002/ar.23559>
6. Tardieu, C., Bonneau, N., Hecquet, J., Boulay, C., Marty, C., Legaye, J., & Duval-Beaupère, G. (2013). How is sagittal balance acquired during bipedal gait acquisition? Comparison of neonatal and adult pelvis in three dimensions. *Evolutionary implications. Journal of Human Evolution*, 65(2), 209-222.
7. Schwab, F., Ungar, B., Blondel, B., et al. (2012). Scoliosis Research Society—Schwab adult spinal deformity classification: a validation study. *Spine*, 37(12), 1077–1082.
8. Stagnara, P., De Mauroy, J. C., Dran, G., et al. (1982). Reciprocal angulation of vertebral bodies in a sagittal plane: approach to references for the evaluation of kyphosis and lordosis. *Spine*, 7(4), 335–342.
9. Roussouly, P., et al. (2005). Classification of the Normal Variation in the Sagittal Alignment of the Human Lumbar Spine and Pelvis in the Standing Position. *Spine*, 30(3), 346-353.
10. Roussouly, P., & Pinheiro-Franco, J. L. (2011). Sagittal parameters of the spine: biomechanical approach. *European Spine Journal*, 20(Suppl. 5), 578-585.
11. Laouissat, F., et al. (2017). Classification of normal sagittal spine alignment: refounding the Roussouly classification. *European Spine Journal*, 27(8), 2002-2011. <https://doi.org/10.1007/s00586-017-4987-9>
12. Borisov, V. N., & Prokhorov, M. A. (1982). Curvature of the human sacrum based on morphometry. *Arkhiv Anatomii Gistologii I Embriologii*, 8, 72–75.
13. Abitbol, M. M. (1989). Sacral curvature and supine posture. *American Journal of Physical Anthropology*, 78(3), 357-366.
14. McKay, G., Torrie, P. A., Dempster, G., et al. (2018). The Relationship between Sacral Kyphosis and Pelvic Incidence. *Asian Spine Journal*, 12(1), 74.
15. Plochocki, J. H. (2011). Sexual Dimorphism of Anterior Sacral Curvature. *Journal of Forensic Sciences*, 56(3), 662-667. doi: 10.1111/j.1556-4029.2010.01455.x
16. Song, K., Wu, B., & Cheng, J. (2020). Correlation study of sacral morphology and sagittal spino-pelvic alignment. *Chinese Journal of Spine and Spinal Cord*, (6), 493-499.
17. Legaye, J., Duval-Beaupère, G., Hecquet, J., & Marty, C. (1998). Pelvic incidence: a fundamental pelvic parameter for three-dimensional regulation of spinal sagittal curves. *European Spine Journal*, 7(2), 99-103.

<https://doi.org/10.1007/s005640050>

17. Schwab, F., Lafage, V., Patel, A., et al. (2009). Sagittal Plane Considerations and the Pelvis in the Adult Patient. *Spine*, 34(17), 1828-1833.
18. Lafage, V., Schwab, F., Patel, A., et al. (2009). Pelvic Tilt and Truncal Inclination: Two Key Radiographic Parameters in the Setting of Adults With Spinal Deformity. *Spine*, 34(17), E599-E606.
19. Barrey, C., Roussouly, P., Le Huec, J. C., et al. (2013). Compensatory mechanisms contributing to keep the sagittal balance of the spine. *European Spine Journal*, 22(6), 834-841.