1	Full title: The landscape and level of alcohol policy enforcement in Tanzania
2	Short title: Alcohol policy enforcement in Tanzania
3	Kim Madundo <sup>a,b</sup> , Aliza Hudda <sup>c</sup> , Maaike L. Seekles <sup>c</sup> , Blandina Mmbaga <sup>d,e</sup> , Angela Obasi <sup>c,f</sup>
4	(a) Department of Mental Health and Psychiatry, Kilimanjaro Christian Medical University
5	College, Moshi, Tanzania
6	(b) Department of Mental Health and Psychiatry, Kilimanjaro Christian Medical Centre, Moshi,
7	Tanzania
8	(c) The Liverpool School of Tropical Medicine, Department of International Public Health,
9	Liverpool, UK
10	(d) Department of Paediatric and Child Health, Kilimanjaro Christian Medical University College,
11	Moshi, Tanzania
12	(e) Kilimanjaro Clinical Research Institute, Moshi, Tanzania
13	(f) AXESS Sexual Health, Royal Liverpool University Hospitals NHS Foundation Trust,
14	Liverpool, UK
15	Corresponding author: Kim Madundo, kimmadundo@yahoo.com
16	
17	
18	
19	
20	
21	

## **Abstract**

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

Harmful use of alcohol causes more deaths in Sub-Saharan Africa than in any other region. In Tanzania, where alcohol use disorders rates are twice the overall African average, harmful alcohol consumption is a public health concern. Given the lack of a contemporary overview of the alcohol policy landscape, we conducted a mixed-methods review of key alcohol-related policies, implementers, and initiatives in Tanzania. We conducted a desk-based review of policy-related documents, and in-depth interviews with eight key informants guided by the 10-compositeindicator framework of a tool for measuring alcohol policy implementation developed by World Health Organization. Representatives were from health-service delivery, community-based organizations, governmental organizations, research, and policymakers whose work is related to alcohol in Tanzania. Data was collected in October 2021, June 2022, and finalized in March 2023. Findings were analyzed using Microsoft Word v2021. Themes were identified, collected, combined, and tabulated. Differences were then resolved by first and second authors. Our findings revealed no single comprehensive national alcohol policy. Pending finalization of a draft policy, various documents and actors govern alcohol production, distribution, licensing, and consumption. Little intersectoral linkage between entities contributes to poor enforcement of these regulations. Regulation is stronger in urban areas, and restrictions more effective on industrial alcohol. However, the majority of consumed alcohol in Tanzania is informally-produced, especially in rural settings. Socio-cultural context plays a key role in alcohol production and consumption, contributing to early-age exposure to alcohol. Alcohol is a growing source of revenue for the Tanzanian government and, therefore, imposing further restrictions is a low priority. There are important policy gaps in various sectors pertaining to alcohol regulation. Our results strongly suggest the need for a comprehensive approach to developing an overarching alcohol policy, with

- 46 involvement of key stakeholders, stronger enforcement, and increased awareness, resources,
- 47 and collaborations.

## Introduction

48

49

50

51

52

53

54

55

56

57

58

59

60

61

62

63

64

65

66

67

68

69

70

The harmful use of alcohol causes approximately 3 million deaths every year worldwide (1). In Sub-Saharan Africa (SSA), alcohol accounts for more deaths than in any other region. The reduction of harmful alcohol use is a particular priority in Tanzania, where alcohol use disorders are twice, and rates of heavy episodic drinking are six times, the Africa Region average (2). Much of this increased risk begins in adolescence (10-19y). Early (aged <15y) alcohol initiation is common, and is more likely to cause dependency issues compared to those who begin drinking above 21 (2). Adults with violence-related injuries are more than six times as likely to have a history of harmful alcohol use (3). Hazardous alcohol consumption is associated with higher odds of a comorbid mental disorder than in the general population (4). Harmful drinking has been linked to insufficient regulations and restrictions on exposure to alcohol-related cues (2,5). There is a need to delay and decrease alcohol use, but a lack of national-level integrated regulations to reinforce this (6). Our study aimed to gain an understanding of alcohol related policy and legislation, and to understand how the enforcement of such policies may affect alcohol use. Published by WHO in 2018, The Global Strategy to reduce the harmful use of alcohol is the most comprehensive international alcohol policy document. This report is endorsed by World Health Organization (WHO) Member States, who commit to periodically provide country-level data on alcohol consumption and related harm, and develop and implement national alcohol policies. The strategy recommends a portfolio of national-level policy options and interventions across ten target areas that could be considered for implementation: Leadership, awareness, and commitment; health services' response; community action; drink-driving policies countermeasures; availability of alcohol; marketing of alcoholic beverages; pricing policies; reducing the negative consequences of drinking and alcohol intoxication; reducing the public

72

73

74

75

76

77

78

79

80

81

82

83

84

85

86

87

88

89

90

91

92

93

94

95

health impact of illicit alcohol and informally produced alcohol; and monitoring and surveillance (2). Through these standard ten markers, this framework serves as a reference point for measurement and comparison of enforcement and implementation across countries worldwide. An action plan (2022-2030) was recently adopted to guide the effective implementation of the strategy (7). Separately, in 2019, the East African Community (EAC) regional policy intended to comprehensively address disparity in policies, programs, and practices, and reduce negative outcomes related to harmful use of alcohol and other drugs (8). Aside from a page in the Global Status Report on Alcohol and Health (2018), there is no recent and detailed overview of what the alcohol policy landscape in Tanzania looks like. In response, this paper presents the findings of an iterative, mixed-methods policy review. It aims to provide an overview of key alcohol-related policies and enforced alcohol control measures, including national-level initiatives and stakeholders. **Materials and Methods** Our review was conducted in two stages: a desk-based review of policy-related documents in Tanzania, and in-depth interviews (IDIs) with key informants. WHO-recommended target areas and corresponding indicators for alcohol-related policy and interventions provided the framework for the searches and IDIs and analysis. National-level key informants, initiatives, and implementers who could offer insight into alcohol-related policies and their enforcement across Tanzania were purposively selected. **Data collection** Firstly, two researchers conducted a desk-based review starting on 17th October 2022, which was then updated in March 2023 to ensure any more recent documents were included. A broad Google Search was conducted, using the following terms: '[Policy OR Law OR Act OR regulation OR legislation] AND [Alcohol OR Liquor OR Brew] AND [Tanzania]'. This returned 80 unique and relevant results.

This open search was followed by targeted searches, informed by WHO target areas and indicators for policy implementation (9,10) and a landscape analysis of alcohol-related policies in Tanzania published in 2012 (11).

Relevant search results were downloaded to a shared folder. Content eligible for inclusion was that describing key policy/actors/plans/strategies related to alcohol consumption in Tanzania, and whereby majority of the content was related to alcohol consumption, and enforced or applicable in Tanzania. Information related to policies and interventions was extracted and a directory of all Tanzanian ministries and special offices with governance over alcohol-related activities was created (*Figure 1*). Websites of these actors were searched for any final relevant sources.

# Figure 1. Organogram of government departments relevant to alcohol-related policies in Tanzania.

Secondly, six IDIs were conducted beginning from 10<sup>th</sup> June 2022, and another two ending on 31<sup>st</sup> March 2023. In addition to perspectives from relevant stakeholders, interviews gathered further information on policies and initiatives related to WHO target areas. Informants were representatives of health-service delivery, community-based organizations (CBOs), governmental organizations, researchers, and policymakers whose work is related to alcohol consumption in Tanzania (see Table 1).

**Table 1. Participant characteristics** 

Participant	Gender	Years of	Private/Public	Scale/Level of
Number		Work	Sector	Operations
		Experience		
#1	Male	19	Private	Regional
#2	Male	12	Public	Regional

#3	Male	20	Private	National
#4	Male	20	Private	Regional
#5	Male	10	Public	National
#6	Male	5	Public	Regional
#7	Male	7	Public	National
#8	Female	4	Private	City

We decided on this number of IDIs based on the overlap in responders' areas of expertise and having reached saturation across our data collection. Topic guide questions were designed to cover all target areas (e.g., relevant policies, health service delivery, efforts to combat harmful alcohol use in the community, schools, and workplace, and availability of alcohol), but were adapted to fit with participant's area of expertise/work (Table 2).

# Table 2. In-depth interview content, participants interviewed and their respective areas of expertise

Composite indicator	IDI participant number(s) and respective areas of			
	expertise			
Leadership, awareness, and	#1 – CBO Representative, Peer support			
commitment	#2 – Social Welfare Officer			
	#3 – CBO Representative, Policy maker			
	#5 – Ministry-level Official, Mental Health Professional,			
	Researcher			
	#6 – Mental Health Professional			

	#8 – Marketing Officer
Health services' response	#2, #3, #6
Community action	#1, #2, #3
	#4 – Researcher, Mental Health Professional
	#7 – Curriculum Developer, Educator
Drink-driving policies and	#2, #5, #8
countermeasures	
Availability of alcohol	#1, #2, #4, #6, #8
Marketing of alcoholic	#1, #5, #7, #8
beverages	
Pricing policies	#2, #5, #6, #8
Reducing the negative	#1, #2, #4, #5, #6, #7
consequences of drinking	
and alcohol intoxication	
Reducing the public health	#1, #2, #4, #5, #6, #7
impact of illicit alcohol and	
informally produced alcohol	
Monitoring and surveillance	#4, #5, #6

IDIs lasted an average of 40 minutes, were conducted in English or Kiswahili, and either in-person or virtually through video communications platform 'Zoom' depending on location and preference of informant. Each interview was given a code number. Participants were reimbursed 10,000 Tanzanian shillings for their participation. No identifying information was collected or recorded during IDIs.

129

130

131

132

133

134

135

136

137

138

139

140

141

142

143

144

145

146

147

148

149

IDI recordings were transferred from voice-recorder to a password-protected computer only accessed by KM and AH, who also transcribed the IDIs. These were then translated to English by the research team, allowing for triangulation of the translations for accuracy. Once complete, transcriptions, translations, and IDI recordings were uploaded to a secure server. **Analysis** Google search results were analyzed by KM and AH; themes were identified, collected and combined in tabulated form. Differences were then resolved by KM and AH, and MS was consulted upon any further disagreements. IDIs were analyzed using Microsoft Word v2021. Quotes were interpreted, color-coded and grouped according to their thematic areas based on WHO target areas, by the first author. A checklist was created indicating the status of policy items in Tanzania; absent, partial restriction, or total restriction. To ensure trustworthiness of our findings we used interviewer triangulation; two researchers worked separately, then discussed and resolved differences. A third researcher monitored quality control of transcripts and codes. All data were triangulated and brought together in one Table, to provide an overview of the current alcohol policy landscape in Tanzania. Ethical approval and consent to participate Clearance to conduct this study was granted by research committees at Kilimanjaro Christian Medical University College (Ref. #2551), the National Institute for Medical Research (Ref. NIMR/HQ/R.8a/Vol.IX/3932, and Liverpool School of Tropical Medicine (Ref: 21-075). Prior to IDIs, informants were sent an information sheet describing the purpose of the study and written informed consent was obtained.

151

152

153

154

155

156

157

158

159

160

161

162

163

164

165

166

167

168

169

170

171

Results Findings from our online searches and IDIs are presented according to the framework of composite indicators provided by WHO. See Table 3 for an overview of the reviewed policyrelated documents, along with their linking components within the WHO framework. Table 3. Relevant policy-related documents Error! Reference source not found. Leadership, awareness and commitment The WHO framework states that member states should exhibit leadership and create awareness around challenges associated with alcohol consumption within their respective countries. They should also demonstrate commitment to controlling such challenges through adequately funded, comprehensive and intersectoral national policies (2). Our searches and IDIs revealed no national alcohol policy or subsequent action plan. Participant #5: "First, we do not have such a [alcohol] policy so far. A national alcohol policy - which was being developed through the bench where I am, but it is a draft. I think - to be made about five to seven years ago. I think it lacked financing" Participant #3: "I don't think if there is a policy that talks about the use of alcohol" Pending a single all-inclusive policy, various documents and government authorities govern activities related to alcohol. There is little intersectoral linkage between these documents and bodies. Participant #5: "There is a need for cooperation in the various enforcement agencies so that so that the implementation can be done well" Imposing further restrictions on the alcohol industry and committing to a national policy was consistently reported as a low priority by respondents.

173

174

175

176

177

178

179

180

181

182

183

184

185

186

187

188

189

190

191

192

193

Participant #5: "There is the EAC regional policy [that talks about preventing, controlling and managing alcohol and drugs in the EAC]. But if it has guidance, if it affects laws, if it has an effect here in Tanzania, I haven't seen it yet" Participant #3: "We have participated in the movement to prepare the National Youth Council. The law was passed five or six years ago but the Council has never been founded. If the council is formed, it will develop youth throughout the country. That is where we talk about violence, alcohol, and other deviant behaviour" Overall, participants' responses indicated that the development of a national alcohol policy is a low priority, particularly because alcohol is a lucrative source of revenue. Participant #1: "I don't think [it is a priority]. Because we all know very well through alcohol, they get revenue" Participant #2: "I have not seen it [alcohol policy] as something that has been given much priority. Because alcohol is everyone's favorite but also... it is the country's economy" Participant #8: "I don't think it is a priority" Participant #6: "So it [alcohol policy] is that the issue is not prioritized? Of course" The government creates and encourages awareness activities on alcohol, often under the category of Non communicable diseases (NCDs) (12). This is done primarily through Ministry for Education, Science, and Technology (MoEST), and the Ministry of Health (MoH). Such awareness campaigns often highlight alcohol as a risk for NCDs. The Tanzania government recently launched the NCD Research Agenda and NCD Strategic Plan version 3 to guide these efforts (13). However, community-based organizations have also played a key role in motivating the government to create awareness and commit to this challenge, as reported by a respondent:

195

196

197

198

199

200

201

202

203

204

205

206

207

208

209

210

211

212

213

214

215

216

Participant #3: "They [several Ministers] saw the importance of my organization to start meeting with students to motivate them and remind them about the contents of the policies that concern them. Those students have not found platforms" Health services' response Member states are required to use a multi-faceted and multi-level approach to create effective alcohol policies; individual-, family-, and community-level interventions are necessary through the health sector as a key entry point for people and families who are at risk of harmful alcohol use. The health sector is well-placed to leverage screening and awareness programs, treatment services, and advocate for community support (2). Our searches identified the NCD Program as responsible for monitoring and reporting issues related to NCDs including health conditions caused by and related to alcohol use. However, there are no known focal national-level individuals or organizations allocated to respond specifically to alcohol issues alone. Participant #3: "In terms of alcohol, there is no board that has been created by the government" Recent literature has shown a missed opportunity for screening at the primary-care level (14). While brief interventions, both psychosocial (education for individuals and families, motivational interviews, referral to support groups) and pharmacological are performed in out- and in-patient settings, these are often unaffordable, and there are no standalone treatment guidelines for alcohol-related issues. Participant #2: "I feel that they do not monitor the issue of alcohol. What they are following is to check his health, maybe with reproductive issues. You find it is difficult to monitor because it is not the main activity there"

218

219

220

221

222

223

224

225

226

227

228

229

230

231

232

233

234

235

236

237

238

239

240

Participant #6: "I can say it is challenging. Many hospitals still do not have psychological services. 70%, they cannot afford. There are others who have to be referred to a hospital [for inpatient care] like Lutindi [more than 300km away from here]" Overall therefore, insufficiency of mental health professionals and low integration of mental health in primary healthcare (15), means that screening for alcohol use is almost exclusively done within specialty-care settings, which is often less affordable and accessible (16). Likewise, the available yet insufficient rehabilitation institutions often go beyond their designated scope of work to compensate for an increasing demand. Community and workplace action WHO recommends that member states and other implementers support and encourage communities to be proactive in reducing harmful alcohol use. WHO further recommends performing rapid assessments to identify priorities, and disparities in knowledge and practice within these communities (2). The Tanzanian government has made some commitment to supporting community action, with limited sustainability. One move is attempting to standardize care for people struggling with use of alcohol and other substances by formulating national guidelines for all sober houses in the country, finalized in 2018. According to these guidelines sober houses should provide supportive and controlled living conditions, as well as psychosocial interventions for people with substance use (and often other related mental health) challenges (17). Due to insufficient rehabilitation programs in Tanzania, sober houses often provide both rehabilitation and transitional services as individuals prepare to re-enter their communities (18). **School-based action** While MoEST has created awareness campaigns against harmful use of substances including alcohol, these happen infrequently, and are particularly rare at the university level, where most students are of legal consumption age. There is a lack of youth awareness and prevention programs, and the necessity for youth engagement in community action.

241

242

243

244

245

246

247

248

249

250

251

252

253

254

255

256

257

258

259

260

261

262

263

Participant #7: "The issue of alcohol is included very little on the curriculum. If we go to colleges, they are very specialized and very rare to teach about alcoholism. Maybe it will happen when we discuss another topic, for example, "If you want to succeed in your career as a professional, things that you need to avoid [include] alcoholism". But there is no topic or sub-topic or course or program" Participant #7: "I'm sure I can't name a page or chapter that lets them know the harm of alcohol. The [Education] policy insists we make school-leavers or university-leavers who can use education to master the environment and their lives. It emphasizes that but not directly" The Tanzania Commission for Universities stipulates higher education institutions create conducive environments for proper conduct among students and staff. Universities then create regulations which serve as technical tools to guide disciplinary action and healthcare assistance for students who present with behavioral changes, including those due to alcohol. Participant #7: "Regulations are created by colleges, but Tanzania Commission for Universities also points out it is appropriate for each institution to have examination regulations and those other disciplinary guidelines"

In an attempt to fill this support gap, various non-governmental organizations (NGOs) offer community-based interventions in Tanzania; Alcoholics Anonymous (AA) is the most widespread CBO across Tanzania, with seven active support groups in different cities (19). Other NGOs have created awareness on alcohol and linked people to care, but frequently as part of a program for Human Immunodeficiency Virus (HIV) care (11).

265

266

267

268

269

270

271

272

273

274

275

276

277

278

279

280

281

282

283

284

285

286

Participant #1: "They [government officials] were very happy and said they didn't know if there is such a thing (Alcoholics Anonymous) here in town, they wanted us to give them our contacts. It's rare to get an invitation directly from the government. I haven't seen them again" **Drink-driving policies and countermeasures** WHO recommends establishing legal limits on blood-alcohol content while driving, the use of sobriety checkpoints, and sanctions for infraction of these regulations. It is further recommended that drivers' licenses be provided on a multiple phase system to ensure sufficient training (2). Tanzania has defined a maximum prescribed blood-alcohol limit for drivers, and requires enforcement of checkpoints for selective blood- and breath-testing for suspected driving under influence of alcohol or actual hazardous driving, and penalties for determined offences (20). A graduated licensing system is currently in place (21). Interviewees consistently highlighted poor integration of enforcement agencies, prioritization of driving offences other than alcohol-related ones, and infrequent utilization of sobriety checkpoints. Participant #2: "They [traffic police] will look at other mistakes, "you have overtaken, speeding, you haven't paid insurance", but they don't say "Let's stop him and check if he is drunk." Participant #5: "The willingness of those enforcement agencies to act... maybe they haven't had direction. It requires relations with other governmental and legal institutions, to be able to show the difference" Availability of alcohol WHO recommends restrictions on production, retailing and distribution, legal age of purchase, and drinking in public spaces as means to achieve this (2).

Various documents comment on regulating availability of alcohol in Tanzania; restrictions on sale of alcohol by time and place are provided, as well as governance on handling of already intoxicated persons. However, there are no restrictions for public environments such as petrol stations or specific events including weddings and concerts (22,23). While an age limit of 18 years is stated for purchase of alcohol both on- and off-premises, interviewees frequently reported traditions from across Tanzania that allow and encourage children to engage in alcohol production, distribution, and consumption, particularly in Northern Tanzania.

**Participant #1**: "At home alcohol was used in every event, recreation, celebration. Boys starting from twelve or fourteen [age] serve elderly men. He asks you to taste a little then you give it to him. It was regarded as culture. It's normal"

Participant #4: "Many have started [using alcohol] at a younger age"

Participants noted that alcohol is highly profitable, and retailers may prioritize income over safety of underage customers.

Participant #2: "Sometimes people don't think about safety of those young people, what they look at is money, how much they earn – "If I stop selling to children, they will run elsewhere, and I may be fired because I have chased away customers"

Participant #6: "Under 18, it's not allowed. The law exists but people do not enforce it"

Another interviewee highlighted the practice of unregulated re-packaging and sale of alcohol in public spaces. This is in contravention with the ban on alcohol sachets (24), due to their affordability and inconspicuous packaging, therefore, also easily accessed by youth.

**Participant #6**: "Strong alcohols being sold at bus stops. They sell them in small glasses ranging from 500, 1000 or 2000 shillings. They also sell to adolescents"

310

311

312

313

314

315

316

317

318

319

320

321

322

323

324

325

326

327

328

329

330

331

Marketing of alcoholic beverages WHO's global strategy recommends the use of legally-binding restrictions on marketing, sponsorships, promotions, and product placement that may directly or indirectly target youth; this includes modern marketing strategies such as social media (2). One of our respondents indicated that alcohol advertisements are often self-regulated by media outlets, however, decisions are heavily influenced by peak engagement times. Participant #8: "These media houses, all they think is money. So, it depends on what time of day they can get the most listeners. For example, on [name of radio program redacted for confidentiality purposes] in the morning, we've had ads [on alcohol] go up" There are partial restrictions on sponsorship of events; local brands are prioritized over foreign ones, particularly for sporting events. Alcohol companies are also not allowed to be the sole sponsor of an event as guided by regulations on fair competition (25). Participant #8: "I don't think they are allowed to sponsor everything. They give a certain percentage of money to invest for like, a marathon, concerts, or football. And to mostly be a Tanzanian brand" There are also partial restrictions on sales promotions; auctioning of alcohol products is permissible with appropriate license and premises (22). However, the Act does not comment on free- or below-cost promotions by retailers. Furthermore, our searches and interviews did not produce results on national-level legally-binding regulations on product placement. **Pricing policies** WHO requires member states to establish a system for adjusting prices of alcohol beverages to account for inflation, excise and duty taxes, with effective implementation and review mechanisms (2).

The Tanzania Revenue Authority (TRA) oversees pricing policies in the country; to instate, evaluate, collect, and adjust excise and duty-paid taxes on industrial alcohol products, thereby controlling prices. TRA enforces laws related to these taxes (26,27). However, estimates show that informally produced alcohol, which is often unregulated, constitutes 86% of all alcohol consumed in Tanzania (23), indicating that TRA has little control over majority of alcohol producers.

Participant #5: "The willingness of those enforcement agencies to act, there may be differences"

Reports reveal the alcohol industry as a growing source of revenue for the government (28). Locally-produced alcohol is also relatively more affordable and easily accessible than industrial alcohol. Furthermore, current pricing policies do not account for disproportionate income levels across different parts of Tanzania (5), and most commonly consumed alcohol products remain affordable even for those without reliable income (29).

Participant #6: "These strong alcohols are still very cheap; it becomes a great influence especially for young people"

## Reducing the negative consequences of intoxication

To mitigate negative consequences associated with drinking alcohol, policies should take a harm-reduction approach; to prevent harm among those most at risk. WHO recommends establishing and enforcing legislation against serving already intoxicated customers, ensuring sufficient server training, and requiring informative labels about alcohol-related harm (2).

Tanzania legally requires health warning labels on alcoholic drinks to promote awareness on age restrictions and drinking in moderation. The Tanzania Bureau of Standards has regulated these inclusions to standardize alcoholic products, as well as other items produced in Tanzania (30).

356

357

358

359

360

361

362

363

364

365

366

367

368

369

370

371

372

373

374

375

376

377

Participant #8: "They include a tagline "you can't sell alcohol to someone who is below 18 years" Furthermore, alcohol manufactured industrially must display alcohol content and ingredients on containers (31), however, there are no requirements for information on contents or additives on labels. In connection to the lack of a definition for a standard drink, there are no legal requirements for displaying number of standard drinks on retail containers. Tanzania also lacks legally-binding requirements for training of persons involved in distribution of alcohol products; recruitment of drink servers and sellers is unregulated (23). Reducing the public health impact of illicit alcohol and informally-produced alcohol WHO recommends developing monitoring systems and imposing the same regulations on informally-produced or unrecorded alcohol as with industrial alcohol. Member states are required to develop monitoring systems for the detection of such alcohol, to issue warnings on the health risks associated with their consumption, and to impose the same regulations on unrecorded alcohol as with industrial alcohol (2). Tanzania has numerous policy-related documents in place that legally require registration of alcohol producers (32), regulation of manufacturing (33), and retail and distribution of alcohol (22,34). However, since more than 80% of consumed alcohol in Tanzania originates from informal producers (23), coupled with low enforcement, the impact of these regulations is likely to be low, and particularly in rural areas. Monitoring and surveillance WHO recommends national surveys on patterns of alcohol consumption and related harms, and establishment of bodies to supervise research and reporting of findings (2).

All IDIs and online searches indicated lack of national-level surveillance of alcohol consumption patterns, health and social consequences, and alcohol policy responses.

WHO has reported data on alcohol consumption levels and trends in Tanzania as part of a larger international survey (23). The STEPwise approach to NCD risk factor Surveillance (STEPS) Survey, also conducted on an international level through WHO, last reported on determinants of NCDs in Tanzania including alcohol in 2012 (35). Furthermore, there has been a rise in research activities investigating determinants and outcomes of alcohol consumption in Tanzania over the past decade (5,11,36,37).

## **Discussion**

This review has revealed valuable insights into policies, laws, acts, and actions regarding alcohol use in Tanzania. There is no national alcohol policy in Tanzania. By using the framework of composite indicators created by WHO, we explored the extent to which alcohol policy is being enforced in the country. Our results indicate a lack of engagement between key sectors, and enforcement, which can have a detrimental effect on effectiveness of existing regulations, and make alcohol more available; this phenomenon has been observed in a previous policy analysis (38). A regional policy scoring scheme was conducted in 46 African countries and revealed Tanzania in 22<sup>nd</sup> place, with medium restrictiveness. This study found a significant association between lower restrictiveness and higher per capita alcohol consumption (39).

While the EAC regional policy document guides prevention programs, treatments services and coordination of policy, its framework is limited in comparison to the WHO Global Strategy; the former has fewer indicators of implementation, and is derived from and influenced by literature on HIV, illicit drugs as well as alcohol. These key differences guided our selection of the WHO document as our reference for the study.

Lack of financing has partly hindered finalization of the national alcohol policy draft. There is minimal intersectoral linkage between various existing regulations and enforcement agencies; minimal to no regulation on marketing of alcohol, pricing policies, and national-level surveillance of consumption trends and socio-economic and health outcomes. Commitment from the government is more impactful on industrial alcohol producers, which accounts for less than 20% of consumed alcohol in the country (5). Furthermore, the majority of our findings indicate that alcohol policy is a relatively low priority for the government, and that an existing regional policy has negligible impact on national-level implementation.

## Interpretation and integration with previous literature

Our research adds to the limited previous literature in this area. For example, a 2012 situational analysis highlighting resources across Tanzania related to alcohol use and its regulation, focused primarily on HIV prevention (11). By contrast, our use of the WHO framework allowed a more comprehensive assessment and comparability with other countries. WHO provides clear guidance on requirements for its member states; Firstly, a primary responsibility for formulating, implementing, monitoring and evaluating public policies to reduce the harmful use of alcohol. The lack of integration and engagement noted in our findings is to be expected given the lack of a single comprehensive policy. This is similarly emphasized in a regional assessment of alcohol policy in African countries, whereby re-evaluation and increasing restrictiveness and integration of policy was recommended (39). Our findings also suggest that through utilizing a socioecological approach, the health sector can support multi-level interventions and advocate for support within communities at large. This is supported by previous literature from Tanzania on missed opportunities to addressing motivators and barriers to seeking care (14).

Stricter regulations on physical access to alcohol are an effective, sustainable, and feasible approach to counteract inappropriate or harmful alcohol use; this is particularly important amongst the young. Marketing restrictions, by extension, are a key point of intervention to minimize early

427

428

429

430

431

432

433

434

435

436

437

438

439

440

441

442

443

444

445

446

447

448

and inappropriate exposure to alcohol advertising. Our findings indicated strong socio-cultural drivers for underage drinking, therefore, strategies to address such practices are paramount. Prospective literature and recent studies from Tanzania similarly indicate that exposure to outdoor advertisements is one of the primary reasons for alcohol initiation among youth (37,40), and highdensity advertising facilitates underage drinking (5). The Tanzanian government has recently made an effort to minimize physical access to alcohol among youth by banning the more affordable alcohol sachets (41). However, our findings indicate that street vendors have found ways to circumvent these restrictions. Raising the cost of alcoholic beverages is one of the most effective strategies for minimizing hazardous alcohol use (42). Our findings revealed that prices of alcoholic beverages remain affordable to most consumers across all ages. Furthermore, unregulated re-packaged alcohol products are available to the public, and are easily accessed by those at risk. Previous literature has explored this gap in policy, and identified pricing regulations as successful in reducing rates of morbidity and mortality (43), violence (44), and unplanned underage pregnancies associated with alcohol use (45). Illegally-imported and informally-produced alcohol (homemade and small-scale production beverages) represent unique subsets of consumed alcohol; their alcohol content may be more difficult to quantify, unknown ingredients may be included, they often cost less than industrial alcohol products, and levels of their consumption are difficult to monitor. Altogether this poses a challenge to member states' efforts in regulating alcohol consumption. Lastly, monitoring and surveillance activities inform enforcement and implementation strategies and allow for more robust policies. Previous studies echo our findings, emphasizing the need for regulation of informal alcohol producers (46) and strengthening monitoring systems (47).

450

451

452

453

454

455

456

457

458

459

460

461

462

463

464

465

466

467

468

469

470

471

472

473

**Strengths and Limitations** To our knowledge, this is the first comprehensive mixed methods alcohol policy review in Tanzania. Additional incorporation of the WHO framework to guide data collection and structuring IDIs adds validity to our findings. It contributes to the previous scholarship on alcohol use in Tanzania, by systematically identifying gaps and opportunities in the alcohol policy landscape in Tanzania through the use of a pre-existing globally recognized framework for measuring policy implementation. Furthermore, by utilizing the WHO framework intended for international audiences, our findings by extension may also offer valuable insights for policy implementation gaps in other countries. We aimed to maximize validity through the rigorous use of multiple methods including a thorough search process, and wide range of IDI participants ranging from primary healthcare delivery, curriculum development, marketing, research, policymaking, CBOs, and Ministry-level officials. These respondents represent settings where mental health and rehabilitation resources for people with a history of hazardous alcohol use are sparse; where people are exposed to production, distribution and advertisements of alcohol from a young age; are known to have little awareness of alcohol and its effects either through education or community programs; and where policy development work has faltered. However, this study had several limitations. The majority of our respondents were representative of regional work and of urban settings, due to challenges in securing interviews with national-level representatives. These findings, therefore, may not provide an entirely accurate representation of enforcement across the country. However, since regulation of alcohol predominantly occurs in urban settings, we are confident in the applicability of these results within that context. Furthermore, several documents were excluded due to their inclusion under HIV programs; frequently, alcohol is discussed as a risk factor for HIV and NCDs. Yet, little has been done to

address socio-cultural complexities around alcohol when included in this way. Pending its own

475

476

477

478

479

480

481

482

483

484

485

486

487

488

489

490

491

492

493

494

495

496

497

498

policy and comprehensive implementation work, alcohol will likely remain in the shadows of other priorities. Undeniably, interviews with individuals regarding aspects of their own professional work may be subject to social desirability biases. However, our desk-review findings corroborate key opinions stated in the IDIs. **Conclusion and recommendations** We have identified lack of overall policy development and co-ordination, poor integration, enforcement and regulation in relation to Alcohol policy in Tanzania. Our findings strongly suggest the need for a comprehensive approach to developing an alcohol policy, with key stakeholders and youth involvement, stronger enforcement, and building on awareness and resources for those affected by and at risk of harmful alcohol use, to address this issue of concern. The health sector has the potential to be a key supporter of awareness activities, mobilizing research around alcohol consumption and its associated challenges, which in turn should inform policy. Further nationwide surveys and other specific studies could add depth to the understanding of alcohol-related harm in Tanzania. The presence and momentum of CBOs and modern marketing strategies are yet another opportunity for low- and middle-income countries such as Tanzania to collaboratively reach, support, and empower people in need. We strongly recommend increasing government support and collaboration with CBOs to improve knowledge on alcohol and standardization of services. Lastly, there is need to publicize and consistently utilize regulations which are already available. List of abbreviations AA - Alcoholics Anonymous CBO - Community-based organization EAC - East African Community HIV - Human Immunodeficiency Virus IDI - In-Depth Interview

499 MoEST - Ministry of Education, Science and Technology 500 MoH - Ministry of Health 501 NCD - Non-Communicate Diseases 502 NGO - Non-governmental organization 503 SSA - Sub-Saharan Africa 504 TRA - Tanzania Revenue Authority WHO - World Health Organization 505 **Acknowledgements** 506 507 Sincere gratitude goes out to the respondents for their time and invaluable experience. 508 **Declarations** 509 **Consent for publication** 510 Not applicable Availability of data and materials 511 512 Deidentified datasets used and/or analysed during the current study are available through URL 513 from the corresponding author on reasonable request. **Competing interests** 514 515 No conflicts of interest were reported by the authors. **Funding** 516 517 This work was supported by a Medical Research Council: Public Health Intervention 518 Development (PHIND) Grant: MR/V032380/1. **Authors' contributions** 519 K.M. led on methodology, study co-ordination, writing, preparation of tables and figures, and 520 521 review & editing. A.H. contributed to methodology, writing, review & editing. M.L.S. contributed 522 to project administration, methodology, study co-ordination, writing, and review & editing. A.O. and B.M. contributed to review & editing. 523

### Patient and Public Involvement

- Patients or the public were involved in the design, or conduct, or reporting, or dissemination
- 526 plans of our research.

### References

524

527

- 1. World Health Organization (WHO). Alcohol. 2022 [cited 2023 Mar 23]. Alcohol. Available from: https://www.who.int/news-room/fact-sheets/detail/alcohol
- 530 2. Hammer JH, Parent MC, Spiker DA, World Health Organization. Global status report on alcohol and health 2018. Vol. 65. 2018. 74–85 p.
- 3. Friedman K, Raymond J, Minnig MC, Sakita F, Vissoci JRN, Staton C. Association between alcohol use and violence-related injury in emergency department patients in Moshi,
  Tanzania: analysis of a prospective trauma registry. Lancet Glob Health. 2020 Apr;8:S18.
- 535 4. Puddephatt JA, Irizar P, Jones A, Gage SH, Goodwin L. Associations of common mental disorder with alcohol use in the adult general population: a systematic review and meta-analysis. Addict Abingdon Engl. 2022 Jun;117(6):1543–72.
- 5. Sommer M, Ibitoye M, Kaaya S, Parker R, Likindikoki S. The Influence of Alcohol Outlet Density and Advertising on Youth Drinking in Urban Tanzania. Health Place [Internet]. 2019;58. Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6708451/
- MOVENDI International. Tanzania: Communities Call for Alcohol Policy to Be Part of
   Election Campaign Discourse [Internet]. 2020 [cited 2023 Mar 23]. Available from:
   https://movendi.ngo/news/2020/10/15/tanzania-communities-call-for-alcohol-policy-to-be-part-of-election-campaign/
- World Health Organization (WHO). Towards an action plan (2022 2030) to effectively implement the Global strategy to reduce the harmful use of alcohol [Internet]. 2022 [cited 2023 Mar 23]. Available from: https://www.who.int/teams/mental-health-and-substance-use/alcohol-drugs-and-addictive-behaviours/alcohol/our-activities/towards-and-action-planon-alcohol
- East African Regional Community. EAST AFRICAN COMMUNITY REGIONAL POLICY ON
   PREVENTION, MANAGEMENT, AND CONTROL OF ALCOHOL, DRUGS AND OTHER
   SUBSTANCE USE. 2019.
- 9. World Health Organization (WHO). Policy in action. A tool for measuring alcohol policy implementation. 2017 p. 1–61.
- 555 10. World Health Organization (WHO). Global strategy to reduce the harmful use of alcohol. Vol. 15. 2016. 1–23 p.
- 11. STRIVE, National Institute of Medical Research (NIMR), Mwanza Intervention Trials Unit
   (MITU). SITUATION ANALYSIS REPORT. HIV PROGRAMMING FOR YOUNG PEOPLE IN
   TANZANIA: EXISTING STRUCTURES AND ALCOHOL RELATED POLICIES. 2012.

12. Ministry of Health, Community Development, Gender, Elderly, and Children (MoHCDGEC).

National Health Policy. Minist Health Community Dev Gend Elder Child Tanzan.

562 2017;(October):1–55.

- 13. United Republic of Tanzania. DRAFT UPDATED MENU OF POLICY OPTIONS AND
   564 COST-EFFECTIVE INTERVENTIONS FOR THE PREVENTION AND CONTROL OF NON 565 COMMUNICABLE DISEASE. 2019 p. 1–4.
- Mushi D, Moshiro C, Hanlon C, Francis JM, Teferra S. Missed opportunity for alcohol use
   disorder screening and management in primary health care facilities in northern rural
   Tanzania: a cross-sectional survey. Subst Abuse Treat Prev Policy. 2022;17(1):1–11.
- 15. Madundo K, Knettel BA, Knippler E, Mbwambo J. Prevalence, severity, and associated
   factors of depression in newly diagnosed people living with HIV in Kilimanjaro, Tanzania: a
   cross-sectional study. BMC Psychiatry. 2023 Feb 1;23(1):83.
- 16. Azevedo MJ. The State of Health System(s) in Africa: Challenges and Opportunities. In:
  Historical Perspectives on the State of Health and Health Systems in Africa, Volume II
- Internet]. Cham: Springer International Publishing; 2017. p. 1–73. Available from:
- 575 http://link.springer.com/10.1007/978-3-319-32564-4 1
- 17. Drug Control and Enforcement Authority (DCEA). Guidelines for the Establishment and Management of Sober Houses in Tanzania Mainland. 2018;1–9.
- 18. Pedersen J, Klepp I. Ten Years After Drug Use and Recovery Among Male Heroin Users in Zanzibar [Internet]. 2021. Available from: http://www.hivolda.no/
- 19. Alcoholics Anonymous. AA Tanzania. 2021. p. 1 Welcome to Alcoholics Anonymous in Tanzania. Available from: https://aatanzania.net
- 582 20. United Republic of Tanzania. CAP. 168 The Road Traffic Act, Amendment of 2019. 2019;1–583 71.
- 21. Tanzania Revenue Authority. Applying driving license for the first time [Internet]. 2023 [cited 2023 Mar 20]. Available from: https://www.tra.go.tz/index.php/2012-11-20-06-37-07/160-applying-driving-license-for-the-first-time
- 587 22. United Republic of Tanzania. The Intoxicating Liquors Act. 1968;1–31.
- 23. World Health Organization (WHO) (WHO). ALCOHOL CONSUMPTION: LEVELS AND
   PATTERNS: Tanzania. World Health Organ 2018. 2016;2016.
- 24. The Citizen. Traders: Alcohol sachets ban big blow. The Citizen [Internet]. 2017; Available from: https://www.thecitizen.co.tz/News/Business/Traders--Alcohol-sachets-ban-big-blow/1840414-3833746-hsxfby/index.html
- 593 25. United Republic of Tanzania. The Fair Competition Act. Parliam United Repub Tanzan. 594 2003;1–38.
- 595 26. United Republic of Tanzania. The Tanzania Revenue Authority Act, 2006. 2006;5–65.

- 596 27. Tanzania Revenue Authority. Tanzania Revenue Aithority: About [Internet]. 2023 [cited 2023 Feb 4]. Available from: https://www.tra.go.tz/index.php/about-tra
- 598 28. AllAfrica. AllAfrica. 2019. p. 1 Tanzania: TRA Records 34% Increased Revenues On Liquor. 599 Available from: https://allafrica.com/stories/201912040349.html
- 29. Francis JM, Weiss HA, Mshana G, Baisley K, Grosskurth H, Kapiga SH. The epidemiology of alcohol use and alcohol use disorders among young people in Northern Tanzania. PLoS ONE. 2015;10(10):1–17.
- 30. Tanzania Bureau of Standards (TBS). Tanzania Bureau of Standards LIQUEURS SPECIFICATIONS. Vol. 10. 2017 p. 0–5.
- 31. United Republic of Tanzania. The Standards Act 2009. 2009;(496):1–41.
- 32. United Republic of Tanzania. The Business Activities Registration Act. 2007;1–11.
- 33. United Republic of Tanzania. The Moshi (Manufacture and Distillation) Act. 1966;(62):1–9.
- 34. United Republic of Tanzania. The Excise (Management and Tariff) Act. 2008;1–76.
- 35. World Health Organization (WHO). TANZANIA STEPS Survey 2012. STEPS Survey. 2012 p. 2–3.
- 611 36. Gerry Mshana. Alcohol taxation, pricing and packaging: experiences from Tanzania and opportunities for structural interventions targeting young people. Mwanza: MITU; NIMR;
- 613 LSHTM; 2014 p. 1–27.
- 37. Mbata D, Mshana G, Francis J, Wamoyi J, Mongi A, Changalucha J, et al. Alcohol Use and Drinking Norms among Young People in Northern Tanzania: Preliminary findings from a qualitative study. STRIVE. 2013.
- 38. Casswell S, Morojele N, Williams PP, Chaiyasong S, Gordon R, Gray-Phillip G, et al. The Alcohol Environment Protocol: A new tool for alcohol policy. Drug Alcohol Rev. 2018;37(November):S18–26.
- 39. Ferreira-Borges C, Esser MB, Dias S, Babor T, Parry CDH. Alcohol control policies in 46 African countries: Opportunities for improvement. Alcohol Alcohol. 2015;50(4):470–6.
- 40. Jernigan D, Noel J, Landon J, Thornton N, Lobstein T. Alcohol marketing and youth alcohol consumption: a systematic review of longitudinal studies published since 2008. Addiction. 2017 Jan;112:7–20.
- 41. The Guardian. Tanzania bans alcohol sachets. The Guardian [Internet]. 2017; Available from: https://guardian.ng/news/tanzania-bans-alcohol-sachets/
- 42. Gibbs N, Angus C, Dixon S, Parry C, Meier P. Effects of minimum unit pricing for alcohol in
   South Africa across different drinker groups and wealth quintiles: a modelling study. BMJ
   Open. 2021 Aug;11(8):e052879.
- 43. Wagenaar AC, Tobler AL, Komro KA. Effects of alcohol tax and price policies on morbidity and mortality: a systematic review. Am J Public Health. 2010 Nov;100(11):2270–8.

- 44. Booth A, Meier P, Shapland J, Wong R, Paisley S. Alcohol pricing and criminal harm: a
   rapid evidence assessment of the published research literature. The University of Sheffield;
   2018 p. 5–6.
- 45. Sen B. Can Beer Taxes Affect Teen Pregnancy? Evidence Based on Teen Abortion Rates and Birth Rates. South Econ J. 2003;70(2):328–43.
- 46. Ferreira-Borges C, Parry CDH, Babor TF. Harmful Use of Alcohol: A Shadow over Sub-Saharan Africa in Need of Workable Solutions. Int J Environ Res Public Health [Internet]. 2017 Mar 27;14(4). Available from: http://www.ncbi.nlm.nih.gov/pubmed/28346373
- 47. Rehm J, Poznyak V. On monitoring unrecorded alcohol consumption. Alcohol Drug Addict. 2015 Jun;28(2):79–89.
- 48. Ministry of Health, Community Development, Gender, Eldery and Children (MoHCDGEC).
   Standard Treatment Guidelines and National Essential Medicines List Tanzania Mainland,
   Sixth Edition, 2021. Gov Tanzan. 2021;93(3):799–801.
- 49. National AIDS Control Programme; MoHCDGEC; Tanzania TUR of. National Guidelines for the Management of HIV and AIDS. Development. 2019;7.
- 50. United Republic of Tanzania. The Law Of The Child Act, 2009. 2009;1–79.
- 51. United Republic of Tanzania. The Public Health Act, 2009. 2009;1–102.
- 52. Tanzania Ministry of Health and Social Welfare (MoHSW); National Institute of Medical
   Research (NIMR); World Health Organization (WHO). Tanzania STEPS Survey Report,
   2013. National Institute for Medical Research (NIMR). 2013 p. 1–154.
- 653 List of figures

- Figure 1. Directory of government actors essential to alcohol-related policies in Tanzania.
- 655 List of tables
- Table 1. Participants' areas of work
- Table 2. Significant statements and relevant policy-related documents
- 658 Table 4. Relevant policy-related documents Error! Reference source not found.

Theme	Sub-theme	Relevant policy-related documents

Leadership,	Written	No national policy
awareness and	national policy	
commitment		The Moshi (Manufacture and Distillation Act (1966)
		(33):
		An Act to regulate the production, sale, and distribution
		of alcohol in Tanzania
		The Business Activities Registration Act (2007) (32): An
		Act to govern registration of all businesses including
		industrial alcohol producers
		The Intoxicating Liquors Act (1968) (22): An Act to
		govern alcohol licensure; legality of premises, hours,
		and environments regarding alcohol; and provides
		definition of an alcoholic beverage
		The Road Traffic Act (2002) (20): Road safety, breath
		testing, sobriety checkpoints
		The Standards Act (2009) (31): An Act that regulates
		standardization of locally-produced and imported
		products; Ensuring quality control
	National	No specific national action plan
	action plan	
		East African Community Regional Policy on Prevention,
		Management, and Control of Alcohol, Drugs, and Other
		Substance Use (2019) (8): A comprehensive policy to
		bridge weaknesses in national policies, programs, and
		other related documents

	Presence of	No national-level awareness-raising activities
	awareness-	specifically targeting alcohol use
	raising	
	activities	
Health services'	Presence of a	No known individual- or organization-based focal points
response	national focal	allocated to respond to alcohol issues alone on a
	point whether	national-level
	organizational	
	or individual,	Standard Treatment Guidelines and National Essential
	for monitoring	Medicines List Tanzania Mainland, Sixth Edition (2021)
	and reporting	(48)
	alcohol-	National Guidelines for the Management of HIV and
	related harm	AIDS, Seventh edition (2019) (49): Contains chapter on
		evidence-based essential treatment for hazardous
		alcohol use
Community and	National	No known national-level support for community action
workplace action	support for	
	community	No known national-level legislature that explicitly
	action	requires school-based alcohol prevention or reduction
		programs
		In development – National guidelines for sober houses
Drink-driving	National	"Eighty milligrams alcohol per hundred milliliters of
policies and	minimum	blood (0.08%)"
countermeasures	legal blood	

	alcohol	-Road Traffic Act (2002) (20): An Act to govern
	concentration	registration, and appropriate and safe use of motor
	(BAC) when	vehicles. Provides definition of legal blood alcohol limit
	driving a	
	vehicle (as a	
	percentage).	
	Sobriety	Road Traffic Act (2002) (20)
	checkpoints	
	Random	Road Traffic Act (2002) (20)
	breath-testing	
	Graduated	Applying driving license for the first time – Tanzania
	licensing	Revenue Authority (21)
Availability of	National	The Moshi (Manufacture and Distillation Act (1966) (33)
alcohol	control of	
	production,	The Excise (Management and Tariff) Act (2008) (34):
	import, sale,	Regulates creation of excisable goods, and collection of
	distribution	excise duties
	and export	
		The Intoxicating Liquors Act (1978) (22)
	National legal	The Intoxicating Liquors Act (1978) (22)
	minimum age	
	for on-/off-	The Law of the Child Act (2009) (50): Consolidates
	premise sales	legislature relating to minors, condemning engagement
	of alcoholic	of children in potentially harmful activities including use
	beverages	of alcohol

	Restrictions	The Intoxicating Liquors Act (1978) (22)
	for on-/off-	
	premise sales	
	of alcoholic	
	beverages	
	Restrictions	The Public Health Act (2009) (51): Regulates conduct
	on drinking in	of alcohol outlets and recreational spaces in the interest
	public places	of promoting, preserving, and maintaining public health
Marketing of	Alcohol	No known legally-binding restrictions on alcohol-
alcoholic	advertising	advertising at the national-level
beverages	Product	No known legally-binding restrictions on product
	placement	placement at the national-level
	Alcohol	The Fair Competition Act (2003) (25): Promotes and
	sponsorship	protects competition in trade, and protects consumers
		from misconduct related to market and commerce
	Sales	The Intoxicating Liquors Act (1978) (22)
	promotions	
Pricing policies	Inflation	Tanzania Revenue Authority Act (2006) (26):
	adjustment	Legislation on pricing policies, excise duties, and
		impartial taxes on behalf of the Central Government
	Excise tax	The Excise (Management and Tariff) Act (2008) (34)
	Duty tax	The Excise (Management and Tariff) Act (2008) (34)
	Price	Tanzania Revenue Authority Act (2006) (26)

	measures	
	other than	
	taxation	
Reducing the	Health	The Standards Act (2009) (31): Governs fees and
negative	warning labels	charges related to products produced in and imported
consequences of		in Tanzania
intoxication	Consumer	The Standards Act (2009) (31)
	information	
	Standard	No known definition of a standard drink
	drinks	
	Alcohol	The Standards Act (2009) (31)
	content	
	Server	No known legally-binding requirements for server
	training	training
Reducing the	Preventing	The Business Activities Registration Act (2007) (32)
public health	illegal	
impact of illicit	production	
alcohol and	Preventing	The Intoxicating Liquors Act (1978) (22)
informally-	illegal sale	
produced alcohol		
Monitoring and	National	National report on consumption levels and trends
surveillance	monitoring	(through WHO) (23): Report prepared by WHO with
	systems	data estimates for Tanzania
	National	STEPS Survey (through WHO) (52): A standardized

surveys	method conducted through WHO for collection,
	analysis, and dissemination of NCD data




Fig 1. Government organogram